

ON VOLUME-PRESERVING VECTOR FIELDS AND FINITE TYPE INVARIANTS OF KNOTS.

R. KOMENDARCZYK AND I. VOLIĆ

ABSTRACT. We consider the general nonvanishing, divergence-free vector fields defined on a domain in 3-space and tangent to its boundary. Based on the theory of finite type invariants, we define a family of invariants for such fields, in the style of Arnold's asymptotic linking number. Our approach is based on the configuration space integrals due to Bott and Taubes.

CONTENTS

1. Introduction	1
Acknowledgments	7
2. Some metric properties of blowups	7
3. Configuration space integrals	8
4. Proofs of Theorems A and B	15
4.1. Key Lemma	16
4.2. Short paths	23
4.3. Proofs of Theorems A, Corollary A, and Theorem B	24
5. Quadratic helicity, energy, and proof of Theorem C.	27
References	29

1. INTRODUCTION

Suppose we have a volume-preserving vector field X defined in some compact domain \mathcal{S} of \mathbb{R}^3 and tangent to its boundary. In the ideal hydrodynamics or magnetohydrodynamics (MHD), c.f. [6] for a comprehensive reference, X plays a role of a vorticity field or a magnetic field. Euler equations (in the ideal hydrodynamics or the ideal MHD) tell us that the flow ϕ_X of X evolves in time under volume-preserving deformations. Therefore, quantities associated

Date: August 26, 2014.

2010 Mathematics Subject Classification. Primary: 57M25; Secondary: 37C50, 76W05, 37C15.

Key words and phrases. volume-preserving fields, asymptotic invariants, configuration space integrals, finite type invariants.

The first author acknowledges the support of DARPA YFA N66001-11-1-4132 and NSF DMS grant #1043009.

The second author acknowledges the support of NSF DMS grant #1205786.

with ϕ_X that are invariant under such deformations are of particular interest to these areas of research.

The best known such invariant is the *helicity of X* , which we will denote by $\mathcal{H}(X)$. It was first discovered by Woltjer in [45]. Its topological nature, i.e. the connection to the linking number of a pair of closed curves in space, was first observed in the work of Moffatt [34] and then fully described by Arnold in [4]. This paper concerns the existence and properties of other invariants of volume-preserving fields derived in the style of Arnold from the finite type (or Vassiliev) invariants of knots and links [41, 10, 3, 44] (see also questions in [5, Problem 1990–16] and [6, p. 176]).

In more detail, and following the general idea of [4], recall that a long piece of an orbit $\mathcal{O}_T(x)$ of a vector field X through $x \in \mathcal{S}$ for time T (or a collection of orbits through different points in \mathcal{S}) can be made into a knot (link) by adding a “short arc” (or as many short arcs as there are orbits) $\sigma(x, y)$ connecting its endpoints, i.e.

$$\bar{\mathcal{O}}_T(x) = \mathcal{O}_T(x) \cup \sigma(x, y), \quad \text{where } y = \mathcal{O}_T(x)(T). \quad (1.1)$$

Thus for any $T > 0$ we obtain a family of knots $\{\bar{\mathcal{O}}_T(x)\}_{x \in \mathcal{S}}$. Now let \mathcal{K} be the space of knots (the set of embeddings of S^1 in \mathbb{R}^3 endowed with the \mathcal{C}^∞ topology) and let

$$\mathcal{F}: \mathcal{K} \longrightarrow \mathbb{R}$$

be a function, typically a knot invariant. This function can be restricted to the family $\{\bar{\mathcal{O}}_T(x)\}_{x \in \mathcal{S}}$, resulting in a function

$$\begin{aligned} \lambda_{\mathcal{S}, T}: \mathcal{S} &\longrightarrow \mathbb{R} \\ x &\longmapsto \mathcal{F}(\bar{\mathcal{O}}_T(x)). \end{aligned}$$

This is a prototype for an invariant of ϕ_X under smooth isotopies via diffeomorphisms isotopic to the identity. In order to produce an actual numerical invariant of ϕ_X , and consequently of X , we need to remove the dependence on short arcs. For that reason, for some $m > 0$ (usually an integer), one considers the limit

$$\mathcal{F}^m(X) = \lim_{T \rightarrow \infty} \int_{\mathcal{S}} \frac{1}{T^m} \lambda_{\mathcal{S}, T}(x) \quad (1.2)$$

We will call $\mathcal{F}^m(X)$ the *asymptotic value of \mathcal{F} along the flow of X (of order m)*. Whenever the order m is specified, we may denote $\mathcal{F}^m(X)$ simply by $\mathcal{F}(X)$. If \mathcal{F} is a knot invariant, this usually gives an invariant of X under volume-preserving deformations. In this case, we will refer to $\mathcal{F}(X)$ as an *asymptotic invariant of X (of order m)*.

Replacing a single orbit $\bar{\mathcal{O}}_T(x)$ by a collection of n orbits $\{\bar{\mathcal{O}}_T(x_1), \dots, \bar{\mathcal{O}}_T(x_n)\}$ at distinct points x_1, \dots, x_n of \mathcal{S} , the above philosophy can be applied to an invariant $\mathcal{F}: \mathcal{L}_n \rightarrow \mathbb{R}$, where \mathcal{L}_n is the space of n -component links (defined and topologized analogously to \mathcal{K}).

Arnold showed in [4] that this technique gives, in the case when \mathcal{F} is the the linking number lk of pairs of orbits $\{\mathcal{O}(x), \mathcal{O}(y)\}$, a well defined invariant $\mathcal{H}(X)$ which equals the

above mentioned Woltjer's helicity. Namely, given a divergence-free field X on \mathcal{S} , we have

$$\mathcal{H}(X) = \int_{\mathcal{S} \times \mathcal{S}} \left(\lim_{T \rightarrow \infty} \frac{1}{T^2} \text{lk}(\bar{\mathcal{O}}_T(x), \bar{\mathcal{O}}_T(y)) \right) \mu(x) \times \mu(y), \quad (1.3)$$

where μ is a volume form on \mathbb{R}^3 , and the function under the integral is a well-defined μ almost everywhere integrable function on \mathcal{S} . Arnold called $\mathcal{H}(X)$ the average *asymptotic linking number* of X and showed that $\mathcal{H}(X)$ is invariant under the volume-preserving deformations of X .

More precisely, let $\text{Vect}(\mathcal{S}, \mu)$ be the Lie algebra of smooth volume-preserving vector fields on $\mathcal{S} \subset \mathbb{R}^3$ equipped with a volume form μ . Consider the action by the group of smooth volume-preserving diffeomorphisms of \mathbb{R}^3 (isotopic to the identity), $\text{Diff}_0(\mathbb{R}^3, \mu)$:

$$\begin{aligned} \text{Diff}_0(\mathbb{R}^3, \mu) \times \text{Vect}(\mathcal{S}, \mu) &\longrightarrow \text{Vect}(g(\mathcal{S}), \mu) \\ (g, X) &\longmapsto g_*X, \end{aligned} \quad (1.4)$$

where g_* stands for the pushforward of the vector field X by the diffeomorphism g . Then invariance under the volume-preserving deformations means the invariance under the above action. In other words,

$$\mathcal{H}(X) = \mathcal{H}(g_*X). \quad (1.5)$$

Remark. Observe that $g_*X(x) = \left. \frac{d}{dt} g \circ \phi_X(t, g^{-1}(x)) \right|_{t=0}$. Thus on the level of flows, the action in (1.4) maps the flow $\phi_X = \phi_X(t, x)$ of X to the flow $g \circ \phi_X \circ g^{-1} = g \circ \phi_X(t, g^{-1}(x))$ of g_*X , i.e.

$$\phi_X \longrightarrow g \circ \phi_X \circ g^{-1}. \quad (1.6)$$

In order to state our main results we first need to provide some general information about finite type invariants, leaving further details for Section 3 (or see, for example, [44] for a more detailed reference). The basic object in the theory of these invariants is a graded algebra (over any ring, but for us, this will be \mathbb{R}) of trivalent diagrams (see Figure 1) which we will denote by \mathcal{D} . The subspace of diagrams of degree n consists of those diagrams with $2n$ vertices and is denoted by \mathcal{D}_n , where $k = k(D)$ vertices are on the circle (*circle vertices*), and $s = s(D)$ vertices are off the circle (*free vertices*). Then \mathcal{D} is the direct sum of \mathcal{D}_n for all $n \geq 1$. For each diagram $D \in \mathcal{D}$, we may construct a function on a knot space \mathcal{K} by means of configuration space integrals, denoted as

$$I_D : \mathcal{K} \longrightarrow \mathbb{R}. \quad (1.7)$$

Details about the map I_D are given in Section 3.

Both \mathcal{D} and its dual, $\mathcal{W} = \mathcal{D}^*$, called the space of *weight systems*, are Hopf algebras. More formally, any $W \in \mathcal{W}$ is a finite linear combination of diagrams in \mathcal{D} . Finite type invariants of knots¹ are indexed by the subspace of *primitive weight systems*, and this is the content of the *fundamental theorem of finite type invariants*, originally due to Kontsevich [29]. An

¹The set up for links is analogous.




FIGURE 1. Examples of trivalent diagrams (without labels or edge orientations). The middle diagram is of degree four, while the other two are of degree three.

alternative proof of this is due to Altschuler and Freidel [3], where the finite type n invariant $V_W: \mathcal{K} \rightarrow \mathbb{R}$ associated with the primitive weight system

$$W = \sum_{D \in TD_n} a_D D \in \mathcal{W}, \quad a_D \in \mathbb{R}, \quad (1.8)$$

is a finite linear combination of functions in (1.7):

$$V_W = \sum_{D \in TD_n} a_D I_D + b I_{D_1}, \quad a_D, b \in \mathbb{R}. \quad (1.9)$$

Here $D_1 = \Theta$, and TD_n denotes the set of trivalent diagrams generating \mathcal{D} . For a more precise statement, see Theorem 3.6. Let us denote the part of the sum W corresponding to diagrams with k vertices on the circle by W^k . Thus if W is a degree n weight system, we have $W = \sum_{k=1}^{2n} W^k$, with the top part of W being W^{2n} ; this corresponds to diagrams all of whose vertices are on the circle (such diagrams are called *chord diagrams*). We can then also clearly write

$$V_W = \sum_{k=1}^{2n} V_{W^k} + V_{D_1}. \quad (1.10)$$

We are now ready to state our main result.

Theorem A. *Let X be a volume-preserving nonvanishing vector field on a compact domain $\mathcal{S} \subset \mathbb{R}^3$, tangent to the boundary. We then have:*

- (i) *For any diagram $D \in \mathcal{D}$ of degree n , the asymptotic value $\mathcal{J}_D^k(X)$, $k = k(D)$ of I_D along the flow of X exists.*
- (ii) *For any invariant V_W of type n , the asymptotic invariant $\mathcal{V}_W(X)$ of order $2n$ exists and equals the asymptotic value $\mathcal{V}_W^{2n}(X)$ of $V_{W^{2n}}$ along the flow X .*
- (iii) *$\mathcal{V}_W(X)$ is invariant under the action by volume preserving diffeomorphisms isotopic to the identity.*

Note that, in part (i), $\mathcal{J}_D^k(X)$ is not necessarily an invariant because I_D is not one. Further, we may consider a situation where $\mathcal{V}_W(X) = \mathcal{V}_W^{2n}(X) = 0$ and see if the lower order averages of V_W exist. For instance, if the asymptotic value $\mathcal{V}_W^{2n-1}(X)$ exists, it may provide a lower order asymptotic invariant of X . Inductively, if $\mathcal{V}_W^j(X) = 0$ for $k < j \leq 2n-1$, we may ask if $\mathcal{V}_W^k(X)$ defines an invariant of a lower order (in the sense of definition following (1.2)). While

we do not answer this question in full generality we obtain the following direct consequence of (i) in Theorem A and (1.10).

Corollary A. *Consider a primitive weight system W and suppose for a given k ($k < n$), we have $W^k \neq 0$. Suppose also that the asymptotic value $\mathcal{V}_W^j(X)$ of W vanishes for every $k < j \leq 2n - 1$ as does the asymptotic value $\mathcal{V}_{W^{k+1}}^k(X)$. Then the asymptotic invariant $\mathcal{V}_W(X)$ of order k exists and equals the asymptotic value $\mathcal{V}_{W^k}^k(X)$ of V_{W^k} along the flow X .*

The meaning of lower order invariants is unclear to us at this point. However, the work in [27, 28] on asymptotic Brunnian links shows one possible setting where they might appear.

A closely related result to Theorem A is proven in [25] by Gambaudo and Ghys who consider a signature invariant $\sigma: \mathcal{K} \rightarrow \mathbb{Z}$ of knots and its asymptotic counterpart for ergodic volume-preserving fields X . In particular, they prove that, in the setting of ergodic fields, the associated *asymptotic signature* $\sigma(X)$ is of order 2 and satisfies

$$\sigma(X) = \frac{1}{2} \mathcal{H}(X). \tag{1.11}$$

An extension of this work on ergodic fields to other knot invariants appears more recently in the work of Baader [7, 8]. In addition, Baader and Marché [9] consider asymptotic finite type invariants. The main result of [9] gives an analog of the identity (1.11) for any asymptotic finite type invariant $\mathcal{V}_W(X)$ of order n whenever X is ergodic and W is degree n . Note that Theorem A shows that $\mathcal{V}_W(X) = \mathcal{V}_W^{2n}(X)$ is well-defined for a general nonvanishing field X (on a domain \mathcal{S} in \mathbb{R}^3), and also indicates a possibility for lower order invariants. Our techniques also lead us to the following counterpart of a result in [9].

Theorem B. *Let μ be the standard volume form on \mathbb{R}^3 and let X be an ergodic μ -preserving nonvanishing vector field on a domain \mathcal{S} . Then there exists a singular differential form $\varpi_{W,2n}$ of degree $4n$ on \mathcal{S}^{2n} , such that*

$$\mathcal{V}_W(X) = c_W (\mathcal{H}(X))^n = \int_{\mathcal{S}^{2n}} \varpi_{W,2n} \wedge \underbrace{(\iota_X \mu \times \cdots \times \iota_X \mu)}_{n \text{ times}}, \tag{1.12}$$

where c_W is a constant independent of X , $\iota_X \mu$ is the contraction of X into the form μ , and $\mathcal{H}(X)$ is the helicity defined in (1.3). Moreover, the lower order invariants (if they exist) are given as follows

$$\mathcal{V}_W^m(X) = \int_{\mathcal{S}^{2m}} \varpi_{W,m} \wedge \underbrace{(\iota_X \mu \times \cdots \times \iota_X \mu)}_{m \text{ times}}.$$

Another avenue we explore here are applications to the *energy-helicity problem* as considered by Arnold in [4] (see also [6]). Define the (magnetic) energy of X by

$$E(X) = \int_{\mathcal{S}} |X|^2 d\mu, \tag{1.13}$$

i.e. as the square of the L^2 -norm of X . Consider the problem of minimizing the energy functional E on the orbit $\mathfrak{o}_X = \{g_* X \mid g \in \text{Diff}_0(\mathbb{R}^3, \mu)\}$ of the action (1.4) through a fixed

vector field X . If \mathfrak{o}_X is an orbit through a general volume-preserving field X there may not be a minimizing (smooth) vector field (c.f. [21]). Can the energy be made arbitrary small? Arnold showed in [4] that

$$E(g^*X) \geq C|\mathcal{H}(X)|, \quad (1.14)$$

for any $g \in \text{Diff}_0(\mathbb{R}^3, \mu)$ and for some positive constant C which depends on the “geometry” (i.e. on a choice of the Riemannian metric on \mathbb{R}^3). Since $\mathcal{H}(X)$ is invariant under the action (1.4), the above inequality gives a lower bound for the magnetic energy of X along the orbit, whenever $\mathcal{H}(X) \neq 0$. Since the bound (1.14) is ineffective for vanishing $\mathcal{H}(X)$, Freedman and He [22] showed a sharper bound for the $L^{3/2}$ -energy² of X in terms of the *asymptotic crossing number*³ $c(X)$ of X :

$$E_{3/2}(X) \geq \left(\frac{16}{\pi}\right)^{1/4} c(X)^{3/4} \geq \left(\frac{16}{\pi}\right)^{1/4} |\mathcal{H}(X)|^{3/4}. \quad (1.15)$$

Asymptotic crossing number is not an invariant under the action (1.4), but it leads to a topological lower bound for fluid knots, i.e. divergence-free vector fields constrained to a tube around a knotted core curve K in 3-space. Namely, denoting by $g(K)$ the genus of K , the following estimate is shown in [22]:

$$E_{3/2}(X) \geq \left(\frac{16}{\pi}\right)^{1/4} (2g(K) - 1)^{3/4} \text{Flux}(X), \quad (1.16)$$

where $\text{Flux}(X)$ is the flux of X through the cross-sectional disk of the tube. In Section 5 of this paper we consider the *quadratic helicity* $\mathcal{H}^2(X)$ (recently proposed by Akhmetiev in [1]). Note that $\mathcal{H}^2(X)$ is well defined, thanks to Theorem A applied to the square of the linking number⁴. Based on the estimate (1.15) we show

Theorem C. *We have*

$$E_{3/2}(X) \geq \left(\frac{16}{\pi}\right)^{1/4} \mathcal{H}^2(X)^{3/8} \geq \left(\frac{16}{\pi}\right)^{1/4} |\mathcal{H}(X)|^{3/4}. \quad (1.17)$$

We end this introduction by saying that our techniques are rather different from [24, 25], where the authors build a “combinatorial model” for an ergodic field, and base their considerations on this model. The configuration space integrals have been used by Cantarella and Parsley in [16] to derive an alternative formula for $\mathcal{H}(X)$ and its “higher dimensional” versions. Considerations of the current paper are measure-theoretic and in the simplest case can be compared to the work of Contreras and Iturriaga on the asymptotic linking number in [18].

Lastly, we wish to indicate that in addition to the results mentioned above, there exists a wealth of approaches to the problem of defining helicity-style invariants of volume-preserving fields, or more generally measurable foliations; see for example papers [2, 42, 40, 19, 31, 35, 26, 30] and references given therein.

²recall that L^2 -energy majorizes the $L^{3/2}$ -energy via the Hölder inequality.

³denoted in [22] by $c(X, X)$.

⁴ lk^2 , which is the simplest finite type 2 invariant of 2-component links

Acknowledgments. We are grateful to Rob Ghrist, Chris Kottke and Paul Melvin for the email correspondence. The first author thanks the organizers of *Entanglement and linking* in Pisa 2011, and in particular Petr Akhmetiev for an interesting conversation during that meeting.

2. SOME METRIC PROPERTIES OF BLOWUPS

Before we review configuration space integrals, in this short section we discuss certain properties of blowups needed for later constructions. Throughout this section, M is a smooth compact manifold with corners. We say that L is a *submanifold* of a smooth compact manifold with corners whenever it is a \mathfrak{p} -submanifold in the sense of [33, Page I.12], which means that local charts come from restriction of the ambient charts to coordinate subspaces. The intersection of two submanifolds N and L is called *clean* if and only if it is transverse and $N \cap L$ is a \mathfrak{p} -submanifold. Recall, following [13] and [39, p. 19],

Definition 2.1. *The blowup of a smooth manifold with corners M along a closed embedded submanifold with corners L is the manifold with boundary $\text{Bl}(M, L)$ that is M with L replaced by those points of the unit normal sphere bundle $S(N(L))$ that are actually the images of paths in M . There is a natural smooth map*

$$\bar{\beta}: \text{Bl}(M, L) \longrightarrow M, \tag{2.1}$$

called the blowdown map, and a partial inverse

$$\beta: M - L \longrightarrow \text{Bl}(M, L) - (\bar{\beta})^{-1}(L), \tag{2.2}$$

called the blowup map.

Given a submanifold N of M such that $N = \text{cl}(N - L)$ (“cl” denoting the closure), we define, following [33, Page V.7], the *lift* of N to $\text{Bl}(M, L)$ as

$$\tilde{N} = \text{cl}(\beta(N - L)).$$

Lifting a vector field on M to $\text{Bl}(M, L)$ amounts to lifting the orbits of the flow (c.f. [33]). Then we have the following natural fact about lifts given as Proposition 5.7.2 in [33, Page V.10], which we paraphrase as

Proposition 2.2. *Suppose submanifolds N and L have a clean intersection in M . Then the lift \tilde{N} in $\text{Bl}(M, L)$ is an embedded submanifold of $\text{Bl}(M, L)$ diffeomorphic to $\text{Bl}(N, N \cap L)$.*

As a next step we equip M with a smooth Riemannian metric g_M and construct a certain smooth metric \tilde{g}_M on $\text{Bl}(M, L)$ which agrees with g outside of a δ -tubular neighborhood⁵ $U_\delta(L)$ of L and turns $U_\delta(L) - L$ into a “cylindrical end” of $\text{Bl}(M, L)$ as in Figure 2. More

⁵I.e. the image of a δ -disk bundle of L under the normal exponential map.

precisely, we define

$$\hat{g}_{\text{Bl}(M,L)} = \begin{cases} dt^2 + g_{\partial U_\delta(L)}; & \text{on } (L \times \mathbb{S}^{k-1}) \times (0, \delta] \cong U_\delta(L) - L, \\ g_M; & \text{outside of } U_\delta(L). \end{cases} \quad (2.3)$$

Here $k = \text{codim}(L)$, t parametrizes $(0, \delta]$ segments in $(L \times \mathbb{S}^{k-1}) \times (0, \delta]$, and $g_{\partial(U_\delta(L))}$ is the restriction of g_M to $\partial(U_\delta(L))$. Since \hat{g}_M may not be smooth along $\partial(U_\delta(L))$, we set $\tilde{g}_{\text{Bl}(M,L)}$ to be obtained by smoothing $\hat{g}_{\text{Bl}(M,L)}$ in the intermediate region $U_{\frac{3}{4}\delta}(L) - U_{\frac{1}{4}\delta}(L)$ (see Figure 2). The above construction will be used later in the case of $C[k; \mathbb{R}^3]$ where \mathbb{R}^3 is considered to have the standard metric.

Next, we indicate a natural estimate which will be very useful in the next section.




FIGURE 2. Illustration of the metric introduced on the blowup of a point in \mathbb{R}^2 .

Lemma 2.3. *Let M be a smooth manifold with corners, L a submanifold of M , and ϖ a smooth m -form on $\text{Bl}(M, L)$. Consider a submanifold N of M whose closure is compact and its lift \tilde{N} to $\text{Bl}(M, L)$. Define*

$$A_{\varpi, \tilde{g}} = \sup_{p \in \tilde{N}} \max_{\substack{v_1, \dots, v_m \in T_p \tilde{N}; \\ |v_i|_{\tilde{g}} = 1}} |\varpi(v_1, \dots, v_m)|. \quad (2.4)$$

Then

$$\left| \int_N \beta^* \varpi \right| = \left| \int_{\tilde{N}} \varpi \right| \leq A_{\varpi, \tilde{g}} \text{vol}(\tilde{N}). \quad (2.5)$$

The proof is clear from definitions since $A_{\varpi, \tilde{g}}$ measures a C^0 -norm of ϖ along \tilde{N} .

3. CONFIGURATION SPACE INTEGRALS

This section contains a brief overview of configuration space integrals (also known as Bott–Taubes integrals). This summary is based on [44] and [39]. We also include some technical results about configuration space integrals that will be needed later. The main result for us is Theorem 3.6. Before we describe configuration space integrals, we briefly review the basic notions from the theory of finite type knot invariants. These invariants have been studied extensively in the last twenty years; for more details, see [41], [10] and [17]. In particular, they are conjectured to *separate* knots.

Let \mathcal{K} be the space of knots, i.e. smooth embeddings of S^1 in \mathbb{R}^3 , with the \mathcal{C}^∞ topology. Any knot invariant $V: \mathcal{K} \rightarrow \mathbb{R}$ can be extended to *singular knots*, which are knots except

for a finite number of transverse self-intersections, using the *Vassiliev skein relation* given in Figure 3. The figure is supposed to indicate that all the singularities have been resolved

$$V\left(\begin{array}{c} \nearrow \\ \bullet \\ \nwarrow \end{array}\right) = V\left(\begin{array}{c} \nearrow \\ \nearrow \\ \nwarrow \end{array}\right) - V\left(\begin{array}{c} \nearrow \\ \nwarrow \\ \nwarrow \end{array}\right)$$

FIGURE 3. Vassiliev skein relation.

(so a knot with n singularities produces 2^n ordinary knots) and V is evaluated on all the resulting knots.

Definition 3.1. *An invariant V is finite type n or Vassiliev of type n if it vanishes on singular knots with $n + 1$ singularities.*

Let \mathcal{V}_n be the real vector space generated by all type n invariants and let $\mathcal{V} = \bigoplus_{n \geq 0} \mathcal{V}_n$. It is immediate that $\mathcal{V}_{n-1} \subset \mathcal{V}_n$, so that one can consider the quotient $\mathcal{V}_n / \mathcal{V}_{n-1}$ (which will appear in Theorem 3.6).

Finite type invariants are intimately connected to the combinatorics of *trivalent diagrams*.

Definition 3.2. *A trivalent diagram D of degree n is a connected graph consisting of an oriented circle, $k = k(D)$ vertices on the circle (*circle vertices*), $s = s(D)$ vertices off the circle (*free vertices*), and some number of edges connecting those vertices. The vertex set $\mathcal{V}(D)$ has cardinality $k + s = 2n$, and all vertices are trivalent (the circle adds two to the valence of a circle vertex), from which it follows that the edge set $\mathcal{E}(D)$ is of cardinality $\frac{k+3s}{2}$. The vertices are labeled by the set $\{1, \dots, 2n\}$, and this labeling induces an orientation on the edges in $\mathcal{E}(D)$ (from the lower-labeled end vertex to the higher-labeled one). We will denote by (i, j) the edge connecting vertices i and j where $i < j$. The diagram is regarded up to orientation-preserving diffeomorphisms of the circle.*

Examples of trivalent diagrams (without labels or edge orientations) are presented in Figure 1. Let TD_n denote the set of trivalent diagrams of degree n and let \mathcal{D}_n be the real vector space generated by TD_n modulo subspaces generated by the *STU relation* illustrated in Figure 4.⁶ Vector space $\mathcal{D} = \bigoplus_{n \geq 0} \mathcal{D}_n$ is in fact a commutative and co-commutative Hopf




FIGURE 4. The *STU* relation: $S = T - U$.

⁶See [44, p. 3] for more details on the *STU* relation.

algebra [10, Theorem 7], where the product (and co-product) is derived from the operation of connected sum of knots. The dual $\mathcal{W} = \mathcal{D}^*$ of \mathcal{D} is known as the space of *weight systems*, with \mathcal{W}_n denoting its degree n subspace, i.e. the dual of \mathcal{D}_n . Since \mathcal{W} also has the structure of a Hopf algebra it is sufficient to understand its primitive elements, called *primitive weight systems*. These generate the entire algebra. A primitive weight system is one that vanishes on *reducible* diagrams, namely those that are not obtained from two diagrams by connected sum (this informally means that, in an irreducible diagram, one cannot draw a line separating $\mathcal{V}(D)$ and $\mathcal{E}(D)$ into two nonempty disjoint subsets).

We now turn our attention to the configuration space integrals. For a manifold M , let $C(q; M)$ be the ordered configuration space of q points in M (i.e. the q -fold product M^q , with the thick diagonal removed). Also recall that, given a submanifold N of a manifold M , the *blowup of M along N* , $\text{Bl}(M, N)$, is obtained by replacing N by the unit normal bundle of N in M (see Definition 2.1). Finally, for S a subset of $\{1, \dots, q\}$, let M^S be the product of $|S|$ copies of M in M^q , indexed by the elements of S , and let Δ_S be the corresponding (thin) diagonal in M^S .

Now let

$$A[k; M] = M^k \times \prod_{S \subset \{1, \dots, k\}, |S| \geq 2} \text{Bl}(M^S, \Delta_S).$$

Definition 3.3. *The Fulton-MacPherson compactification of $C(k; M)$, denoted by $C[k; M]$, is the closure of the image of the inclusion*

$$\alpha_M : C(k; M) \longrightarrow A[k; M], \tag{3.1}$$

where the S -factors of this map are given by the blowup maps⁷. We denote α_M by α if M is understood, and we will also refer to it as the blowup map of $C(k; M)$. The blowdown map $\bar{\alpha}_M : C[k; M] \longrightarrow M^k$ is obtained by the obvious restriction of the projection of $A[k; M]$ onto its M^k factor.

Equivalently, $C[k; M]$ can be obtained from M^k by successive blowups of Δ_S diagonals in M^k [13, 39]. These blowups have to be performed in the order dictated by the inclusion relation \subset on the indexing sets S . More precisely, if $S' \subset S$, then $\Delta_{S'}$ should be blown up before Δ_S . Yet another equivalent definition is due to Sinha [37]. All these definitions produce diffeomorphic smooth manifolds with corners, compact when M is compact, and homeomorphic to a complement of a tubular neighborhood of the thick diagonal in M^k . The interior of $C[k; M]$ equals the image of $C(k; M)$ under α and will be denoted by $C_0(k; M)$. For the remainder of this section we will mostly need the case $M = \mathbb{R}^3$. In this situation, one needs to equip the compactification $C[k; \mathbb{R}^3]$ with a face at infinity for it to be a compact manifold with corners. We also point out that compactification is functorial and in particular we have

⁷see Equation (2.2)

Proposition 3.4 ([23, 37]). *Suppose $g : M \rightarrow N$ is an embedding of a smooth manifold M into a smooth manifold N . We then have an induced embedding*

$$\tilde{g} : C[k; M] \longrightarrow C[k; N]$$

of manifolds with corners, which respects the boundary stratifications and extends the obvious product map $g^k : C(k; M) \longrightarrow C(k; N)$, $g^k = g \times \cdots \times g$, such that the following diagram commutes

$$\begin{array}{ccc} C[k; M] & \xrightarrow{\tilde{g}} & C[k; N] \\ \uparrow \alpha_M & & \uparrow \alpha_N \\ C(k; M) & \xrightarrow{g^k} & C(k; N). \end{array} \quad (3.2)$$

The reader may consult, for example, [37, Corollary 4.8] for a proof of this proposition.

Given the compactified configuration space $C[g; \mathbb{R}^3]$ and any two positive integers k and s , define $C[k, s; \mathcal{K}, \mathbb{R}^3]$ to be the pullback bundle in the following diagram

$$\begin{array}{ccc} C[k, s; \mathcal{K}, \mathbb{R}^3] & \xrightarrow{p_{k,s}} & C[k + s; \mathbb{R}^3] \\ \downarrow \bar{\pi}_k & & \downarrow \pi_k \\ C[k; S^1] \times \mathcal{K} & \xrightarrow{\tilde{\text{ev}}} & C[k; \mathbb{R}^3], \end{array} \quad (3.3)$$

where π_k is the usual projection onto the first k coordinates and

$$\tilde{\text{ev}}(\cdot, K) : C[k; S^1] \longrightarrow C[k; \mathbb{R}^3]$$

is the evaluation map induced from the knot embedding map $K : S^1 \hookrightarrow \mathbb{R}^3$; see Proposition 3.4. In other words it is a “lift” of the product map

$$\begin{aligned} \text{ev} : C(k; S^1) \times \mathcal{K} &\longrightarrow C(k; \mathbb{R}^3) \\ ((t_1, \dots, t_k), K) &\longmapsto (K(t_1), \dots, K(t_k)) \end{aligned} \quad (3.4)$$

to the compactified spaces. All maps in Diagram (3.3) are smooth maps of manifolds with corners [13, 37], which is equivalent to saying that they admit smooth extensions to some open neighborhoods of the domains of their charts.

Returning now to the diagram algebra \mathcal{D} , for a trivalent diagram $D \in \mathcal{D}_n$, define the associated Gauss map to be the product

$$h_D = \prod_{(i,j) \in \mathcal{E}(D)} h_{i,j} : C[k, s; \mathcal{K}, \mathbb{R}^3] \longrightarrow \prod_{(i,j) \in \mathcal{E}(D)} S^2, \quad (3.5)$$

where $h_{i,j}: C[k, s; \mathcal{K}, \mathbb{R}^3] \longrightarrow S^2$ is the lift to the compactification of the classical Gauss map

$$\begin{aligned} C(k + s; \mathbb{R}^3) &\longrightarrow S^2, \\ (x_1, \dots, x_i, \dots, x_j, \dots, x_{k+s}) &\longmapsto \frac{x_j - x_i}{|x_j - x_i|}. \end{aligned}$$

Maps $h_{i,j}$ extend smoothly to the boundary of $C[k, s; \mathcal{K}, \mathbb{R}^3]$, [13, Appendix]. Thus h_D is also smooth, and as a result we obtain a smooth $(k + 3s)$ -form ω_D on $C[k, s; \mathcal{K}, \mathbb{R}^3]$ via the pullback:

$$\omega_D = h_D^*(\omega \times \cdots \times \omega) = \prod_{(i,j) \in \mathcal{E}(D)} \omega_{i,j}, \quad \omega_{i,j} = h_{i,j}^* \omega. \quad (3.6)$$

Here ω is the area form on S^2 , usually chosen in standard coordinates on \mathbb{R}^3 as

$$\omega(x, y, z) = \frac{x dy \wedge dz - y dx \wedge dz + z dx \wedge dy}{(x^2 + y^2 + z^2)^{\frac{3}{2}}}.$$

One now has a smooth bundle of manifolds with corners,

$$p_{\mathcal{K}} : C[k, s; \mathcal{K}, \mathbb{R}^3] \longrightarrow \mathcal{K},$$

which is the composition of $\bar{\pi}_k$ with the trivial projection of $C[k; S^1] \times \mathcal{K}$ onto the second factor. The fiber of $p_{\mathcal{K}}$ over a knot K is the configuration space of $k + s$ points in \mathbb{R}^3 , first k of which are constrained to lie on K . Integration along the $(k + 3s)$ dimensional fiber of $p_{\mathcal{K}}$ produces a 0-form (a function) on \mathcal{K} . We will denote its value at $K \in \mathcal{K}$ by $I_D(K)$. In other words,

$$I_D(K) := ((p_{\mathcal{K}})_* \omega_D)(K). \quad (3.7)$$

Remark 3.5. Note that ω_D vanishes to the order $1/r^n$ at “infinity” of $C[k + s; \mathbb{R}^3]$, where r is the distance from the origin. It is therefore integrable along fibers of $p_{\mathcal{K}}$ and thus $(p_{\mathcal{K}})_* \omega_D$ is well-defined.

We now have the following fundamental result originally due to Altschuler and Freidel [3], but reproved by Thurston [39] in the form we use here.

Theorem 3.6 ([3, 39]). *Given a primitive weight system $W \in \mathcal{W}_n$, $n \geq 0$, the map defined by*

$$\begin{aligned} V_W : \mathcal{K} &\longrightarrow \mathbb{R} \\ K &\longmapsto \frac{1}{(2n)!} \sum_{D \in TD_n} W(D) (I_D(K) - m_D I_{\Theta}(K)), \end{aligned} \quad (3.8)$$

where m_D is a real number which depends only on D , is a finite type n knot invariant. Moreover, any finite type invariant of type n can be expressed as V_W for some primitive weight system $W \in \mathcal{W}_n$. More precisely, V_W gives an isomorphism $\mathcal{V}_n / \mathcal{V}_{n-1} \cong \mathcal{W}_n$ for all $n \geq 0$ (where by \mathcal{V}_{-1} we mean the one-dimensional space of constant invariants).

Notice that the statement above is a more elaborate version of (1.9), with $a = \frac{1}{2n!}W(D)$ and $b = \frac{1}{2n!}m_D$. The term $m_D I_{\Theta}(K)$ is known as the *anomalous correction*. The integral $I_{\Theta}(K)$ computes the *writhing number*⁸ of K (see [14]).

We next wish to clarify some technical aspects of the integration in (3.7) that will be needed for later constructions. Let $K \in \mathcal{K}$ and $D \in \mathcal{D}_n$. The Gauss map h_D from (3.5) factors as $h_D = \bar{h}_D \circ p_{k,s}$ (see Diagram (3.3)), where

$$\bar{h}_D : C[k+s, \mathbb{R}^3] \longrightarrow \prod_{(i,j) \in \mathcal{E}(D)} S^2$$

has an identical definition as h_D in (3.5). We also have the analog of the form ω_D on $C[k+s, \mathbb{R}^3]$, given as $\bar{h}_D^*(\omega \times \cdots \times \omega)$. We will also denote this form by ω_D . Integrating along the $3s$ -dimensional fiber of π_k in Diagram (3.3), we obtain a smooth k -form (see [13, p. 5281])

$$\varpi_D = (\pi_k)_* \omega_D \quad (3.9)$$

on $C[k; \mathbb{R}^3]$ (see Remark 3.5). On the other hand, the evaluation map (3.4), produces at each point

$$\text{ev}((t_1, \dots, t_k), K) = (K(t_1), \dots, K(t_k)) \in C(k; \mathbb{R}^3), \quad (t_1, \dots, t_k) \in C(k; S^1),$$

a frame

$$\dot{K}_k = \{\dot{K}(t_1), \dots, \dot{K}(t_k)\}, \quad \dot{K}(t_i) = \frac{d}{dt_i} K(t_i).$$

Lifting this frame to $C[k; \mathbb{R}^3]$, via the map pushforward α_* induced by $\alpha = \alpha_{\mathbb{R}^3}$ from (3.1), we obtain the frame $\tilde{K}_k = \alpha_* \dot{K}_k$. Contracting into ϖ_D , given by (3.9), we obtain a (distributional) function over $C[k; \mathbb{R}^3]$, determined by

$$f_{D,K}(\mathbf{t}) := \varpi_D(\tilde{K}_k)(\mathbf{t}) = \alpha^* \varpi_D(\dot{K}_k)(\mathbf{t}) = (\text{ev}_K^* \alpha^* \varpi_D)(\mathbf{t})[\partial_{\mathbf{t}}], \quad (3.10)$$

for $\mathbf{t} = (t_1, \dots, t_k)$. Here α^* denotes the pullback induced by α .

Proposition 3.7. *With $f_{D,K}$ as defined in (3.10), we have the following identity for $I_D(K)$:*

$$I_D(K) = \underbrace{\int_0^T \cdots \int_0^T}_{k \text{ times}} f_{D,K}(\mathbf{t}) \, d\mathbf{t}, \quad (3.11)$$

where the interval $[0, T]$ parametrizes the knot K .

Proof. Restricting $\bar{\pi}_k$ in Diagram (3.3) to the fiber over the point $K \in \mathcal{K}$ and the rest of the maps to the subset of the interior of the compactifications $C_0(\cdot; \mathbb{R}^3) \subset C[\cdot; \mathbb{R}^3]$, we have a

⁸Note that the writhing number is an average *writhe* over all possible projections of the knot.

diagram

$$\begin{array}{ccccc}
C_0(k, s; K, \mathbb{R}^3) & \xhookrightarrow{j} & C_0(k; S^1) \times C_0(k+s; \mathbb{R}^3) & \xrightarrow{v} & C_0(k+s; \mathbb{R}^3) \\
& \searrow^{\tilde{\pi}_k} & \downarrow & & \downarrow^{\pi_k} \\
& & C_0(k; S^1) & \xrightarrow{\text{ev}_K} & C_0(k; \mathbb{R}^3),
\end{array}$$

where $p = v \circ j$. Since $C(k+s; \mathbb{R}^3) \subset (\mathbb{R}^3)^k \times (\mathbb{R}^3)^s$, we may work with the standard coordinates $(\mathbf{x}, \mathbf{y}) = (x_1, \dots, x_k, y_1, \dots, y_s)$ on $(\mathbb{R}^3)^k \times (\mathbb{R}^3)^s$ imposed by the blowup map $\alpha : C(k+s; \mathbb{R}^3) \rightarrow C_0(k+s; \mathbb{R}^3) \subset C[k+s; \mathbb{R}^3]$ of (3.1). Let components of vectors x_j and y_j in (\mathbf{x}, \mathbf{y}) be further indexed as $x_j = (x_j^i)_{i=1,2,3}$, $y_j = (y_j^i)_{i=1,2,3}$ and denote by $\mathbf{t} = (t_1, \dots, t_k)$ the coordinates on $C(k; S^1) \subset (S^1)^k$. Then the $(k+3s)$ -form $\alpha^* \omega_D$ can be written as

$$\alpha^* \omega_D = d\mathbf{y} \wedge \hat{\alpha}_D + \hat{\alpha}_D,$$

where $d\mathbf{y}$ is the top degree form on $(\mathbb{R}^3)^s$ and $\hat{\alpha}_D$ some $(k+3s)$ -form not containing the term $d\mathbf{y}$. Using the multi-index notation, let us write

$$\hat{\alpha}_D = \sum_{I, J} \hat{a}_{I, J}(\mathbf{x}, \mathbf{y}) d\mathbf{x}_I^J.$$

Here $d\mathbf{x}_I^J = dx_{j_1}^{i_1} \wedge \dots \wedge dx_{j_k}^{i_k}$ and I, J are appropriate multiindices. After integrating along the fiber, we get

$$(\pi_k)_* \alpha^* \omega_D = \alpha^* \varpi_D(\mathbf{x}) = \sum_{I, J} \left(\int_{\pi_k^{-1}(\mathbf{x})} \hat{a}_{I, J}(\mathbf{x}, \mathbf{y}) d\mathbf{y} \right) d\mathbf{x}_I^J,$$

where $\pi_k^{-1}(\mathbf{x}) = C(s, \mathbb{R}^3 - \{x_1, \dots, x_k\})$.⁹ From (3.10), we have

$$f_{D, K}(\mathbf{t}) = \sum_{I, J} \left(\int_{\pi_k^{-1}(K(\mathbf{t}))} \hat{a}_{I, J}(K(t_1), \dots, K(t_k), \mathbf{y}) d\mathbf{y} \right) b_I^J(t_1, \dots, t_k),$$

where $b_I^J(t_1, \dots, t_k) = d\mathbf{x}_I^J[\dot{K}^{\wedge k}]$ and

$$(\text{ev}_K^* \alpha_* \varpi_D)(\mathbf{t}) = f_{D, K}(\mathbf{t}) d\mathbf{t}, \quad d\mathbf{t} = dt_1 \wedge \dots \wedge dt_k.$$

On the other hand, $v^* \alpha^* \omega_D$ has the identical expression as $\alpha^* \omega_D$, so we may write $j^* \alpha^* \omega_D$ for $j^* v^* \alpha^* \omega_D$. In the $(\mathbf{t}, \mathbf{x}, \mathbf{y})$ coordinates, we obtain

$$j^* \alpha^* \omega_D = \sum_{I, J} \hat{a}_{I, J}(K(t_1), \dots, K(t_k), \mathbf{y}) b_I^J(t_1, \dots, t_k) d\mathbf{y} \wedge dt_1 \wedge \dots \wedge dt_k.$$

⁹I.e. $\pi_k^{-1}(\mathbf{x})$ is a configuration space of s points in \mathbb{R}^3 with k points deleted, see [20].

Since the boundary of the configuration space $C[k; S^1]$ is measure zero, it does not contribute to the integral in (3.7) and we easily see that the following identities, proving (3.11), hold:

$$\begin{aligned} I_D(K) &= ((p_X)_* \alpha^* \omega_D)(K) = \int_{C(k; S^1)} (\bar{\pi}_k)_* (\alpha^* \omega_D) \\ &= \int_{C(k; S^1)} (\bar{\pi}_k)_* (j^* \alpha^* \omega_D) = \int_{C(k; S^1)} f_{D, K}(\mathbf{t}) \, d\mathbf{t}. \quad \square \end{aligned}$$

4. PROOFS OF THEOREMS A AND B

In the setting of a volume-preserving vector field X on a domain \mathcal{S} from Theorem A, we wish to apply constructions of Section 3 to the family of knots $\{\bar{\mathcal{O}}_T(x)\}$ obtained from the “closed up” orbits of X . Note that any such orbit $\bar{\mathcal{O}}_T(x)$ (as in (1.1)) is generically a piecewise smooth knot in \mathbb{R}^3 . In order to define $\bar{\mathcal{O}}_T(x)$, one needs a system $\{\sigma(x, y)\}$ of *short paths* on \mathcal{S} , which can in general be defined from geodesics after an appropriate choice of the metric on \mathcal{S} [43]. Short paths will in particular be dealt with in Lemma 4.5. The main property of short paths we will use is that their length is uniformly bounded. Note that we can assume $\bar{\mathcal{O}}_T(x)$ is smooth, because its corners can be rounded and $\bar{\mathcal{O}}_T(x)$ is the C^0 limit of these “rounded” parametrizations.

Recall that the basic ingredient of the formula (3.8) for any finite type n invariant V_W is the integration function I_D associated with a diagram $D \in \mathcal{D}_n$. Following the ideas outlined in the Introduction we focus on the family of functions

$$\begin{aligned} \mathcal{S} &\longrightarrow \mathbb{R}, \\ x &\longmapsto I_D(\bar{\mathcal{O}}_T(x)) \end{aligned}$$

that is dependent on T . For any $x \in \mathcal{S}$, we wish to study the time average

$$\bar{\lambda}_D(x) = \lim_{T \rightarrow \infty} \frac{1}{T^k} I_D(\bar{\mathcal{O}}_T(x)), \quad k = k(D). \quad (4.1)$$

Naturally, we need to investigate if $\bar{\lambda}_D$ is a well-defined function on \mathcal{S} and whether it is integrable.

Recall that $X(x) = \dot{\mathcal{O}}_T(x)$. Given a smooth k -form ϖ_D on $C[k; \mathbb{R}^3]$ as defined in (3.9), we have a global analog of the function $f_{D, K}$ in (3.10):

$$f_{D, X} : C(k; \mathcal{S}) \longrightarrow \mathbb{R}, \quad f_{D, X} := \alpha^* \varpi_D(X, \dots, X), \quad (4.2)$$

where the frame of fields $\{X, \dots, X\}$ spans the tangent space to the product of orbits $\mathcal{O}(x_1) \times \dots \times \mathcal{O}(x_k)$ through any point (x_1, \dots, x_k) in \mathcal{S}^k . It is convenient to think about the above constructions in terms of the underlying foliation \mathcal{F}_X^k of \mathcal{S}^k defined by the orbits of the action of the k -fold product flow ϕ_X^k on \mathcal{S}^k . Note that \mathcal{F}_X^k has complete leaves because X is tangent to $\partial\mathcal{S}$, and orbits $\mathcal{O}(x)$ thus exist for all time. The function $f_{D, X}$ is well-defined on $C(k; \mathcal{S})$, but, except for along the orbits, it generally blows up close to the diagonals of \mathcal{S}^k . We can

also consider the function

$$\tilde{f}_{D,X} : C_0(k; \mathcal{S}) \longrightarrow \mathbb{R}, \quad f_{D,X} := \varpi_D(\tilde{X}, \dots, \tilde{X}), \quad (4.3)$$

where $\{\tilde{X}, \dots, \tilde{X}\}$ is a lift of the frame $\{X, \dots, X\}$ of vector fields to $C[k; \mathcal{S}]$. Note that even though ϖ_D is smooth on $C[k; \mathcal{S}]$, the vector field $\tilde{X} = \alpha_* X$ undergoes “infinite stretching” close to the boundary of $C[k; \mathcal{S}]$ (see Remark 4.3). Clearly, $f_{D,X}$ factors as

$$f_{D,X} = \tilde{f}_{D,X} \circ \alpha.$$

Since $\bar{\mathcal{O}}_T(x)$ is parametrized by the interval $[0, T + 1]$ (where $[T, T + 1]$ parametrizes a short segment $\sigma(x, \phi_X(x, T))$), Proposition 3.7 applied to (4.1) pointwise yields

$$I_D(\bar{\mathcal{O}}_T(x)) = \underbrace{\int_0^{T+1} \cdots \int_0^{T+1}}_{k \text{ times}} f_{D,X}((\phi \cup \sigma)(x, t_1), \dots, (\phi \cup \sigma)(x, t_k)) dt_1 \cdots dt_k, \quad (4.4)$$

$$\bar{\lambda}_D(x) = \lim_{T \rightarrow \infty} \frac{1}{T^k} I_D(\bar{\mathcal{O}}_T(x)).$$

Here $(\phi \cup \sigma)$ is a shorthand notation for the flow ϕ of X followed by the short path parametrization.

This is thus the setup in which we will prove our main theorems in this section. But before we can do that, we will establish a useful lemma.

4.1. Key Lemma. Here is the lemma that will be used in the proofs of Lemma 4.5 and Theorems A and B.

Key Lemma. *Let μ be the underlying measure on the domain $\mathcal{S} \subset \mathbb{R}^3$, invariant under the flow of X . Consider the time average of $f_{D,X}$ over \mathcal{F}_X^k , defined as*

$$\lambda_D(x) = \lim_{T \rightarrow \infty} \frac{1}{T^k} \int_0^T \cdots \int_0^T f_{D,X}(\phi(x, t_1), \dots, \phi(x, t_k)) dt_1 \cdots dt_k, \quad x \in \mathcal{S}, \quad (4.5)$$

where in comparison to (4.1), we skipped the integrals over short paths. Then this limit exists almost everywhere on \mathcal{S} and λ_D is in $L^1(\mathcal{S}, \mu)$.

Before we prove this, we need to make several observations. Note that μ induces a measure on \mathcal{S}^k by the pushforward via the thin diagonal inclusion $\mathcal{S} \hookrightarrow \mathcal{S}^k$, $x \mapsto (x, \dots, x)$. Let us denote the resulting measure by μ_Δ . Clearly μ_Δ is a finite Borel measure supported on the thin diagonal of \mathcal{S}^k . Averaging over the \mathbb{R}^k -action of $\phi^k = \phi_X^k$ we obtain a ϕ^k -invariant measure

$$\bar{\mu}_\Delta = \lim_{T \rightarrow \infty} \frac{1}{T^k} \int_0^T \cdots \int_0^T ((\phi^k)_* \mu_\Delta) dt_1 \cdots dt_k. \quad (4.6)$$

For the k -fold product \mathcal{S}^k , the above is a well defined limit in the space of Borel measures $\mathcal{M}(\mathcal{S}^k)$, c.f. [12]. Note that $\bar{\mu}_\Delta$ is supported on the set of leaves of the foliation \mathcal{F}_X^k intersecting

the thin diagonal in \mathcal{S}^k . From the definitions, we may write $\int_{\mathcal{S}} \lambda_D d\mu$ as

$$\begin{aligned} \int_{\mathcal{S}} \lambda_D d\mu &= \int_{\mathcal{S}^k} \left(\lim_{T \rightarrow \infty} \frac{1}{T^k} \int_0^T \cdots \int_0^T f_{D,X}(\phi(x_1, t_1), \dots, \phi(x_k, t_k)) dt_1 \cdots dt_k \right) d\mu_{\Delta} \\ &= \int_{\mathcal{S}^k} f_{D,X} d\bar{\mu}_{\Delta}, \end{aligned} \quad (4.7)$$

where in the third identity we used (4.6). Therefore the question of whether λ_D is in $L^1(\mathcal{S}, \mu)$ is equivalent to the question of whether $f_{D,X}$ of (4.2) is in $L^1(\mathcal{S}^k, \bar{\mu}_{\Delta})$.

Remark 4.1. In place of $\bar{\mu}_{\Delta}$ one can consider any other invariant measure, in particular we may restrict just to any measure supported on the k -product $\mathcal{O}(x) \times \cdots \times \mathcal{O}(x)$ of a single long orbit, or equivalently obtained by averaging, as in (4.6), a Dirac delta measure of a point $(x, \dots, x) \in \mathcal{S}^k$. It is well known (c.f. [12, 15]) that $\bar{\mu}_{\Delta}$ can be arbitrarily well approximated by finite sums of such Dirac delta averages. (We will use this fact in Section 5.)

In order to investigate integrability of $f_{D,X}$, we employ the following natural generalization of [15, Proposition 10.3.2] to a product of flows.

Proposition 4.2 ([15]). *Any ϕ^k -invariant measure μ on \mathcal{S}^k corresponds to a holonomy invariant measure of the foliation \mathcal{F}_X^k .*

Let us choose a finite regular foliated atlas for \mathcal{F}_X^k where a domain V^{α} , $\alpha = (\alpha_1, \dots, \alpha_k)$, of each chart is a product of regular flow boxes $\{V_{\alpha}\}$ of the vector field X covering \mathcal{S} . In other words,

$$V^{\alpha} = V_{\alpha_1} \times \cdots \times V_{\alpha_k}, \quad V_{\alpha_i} = \mathcal{J}_{\alpha_i} \times I_{\alpha_i}, \quad I_{\alpha_i} = (-\epsilon_{\alpha_i}, \epsilon_{\alpha_i}), \quad 0 < \epsilon_{\alpha_i} < \epsilon, \quad (4.8)$$

where each \mathcal{J}_{α_i} is a transverse disk to the flow of X . V^{α} can be expressed as the product

$$V^{\alpha} = \mathcal{J}^{\alpha} \times I^{\alpha} = \left(\prod_i \mathcal{J}_{\alpha_i} \right) \times \left(\prod_i I_{\alpha_i} \right).$$

Recall from [15] that a holonomy invariant measure $\nu_{\mathcal{F}}$ of $\mathcal{F} = \mathcal{F}_X^k$ is a measure defined on $\bigsqcup_{\alpha} \mathcal{J}^{\alpha}$ that is invariant under the action of the holonomy pseudogroup of \mathcal{F} . The Ruelle–Sullivan Theorem [36] (see also [15, p. 245]) and Proposition 4.2 imply the existence of a holonomy invariant (finite) measure $\nu_{\mathcal{F}}$ corresponding to $\bar{\mu}_{\Delta}$, given in (4.6), such that

$$\int_{\mathcal{S}^k} f_{D,X} \bar{\mu}_{\Delta} = \sum_{\alpha} \int_{\mathcal{J}^{\alpha}} \left(\int_{I^{\alpha}} \xi_{\alpha}(\mathbf{x}, \mathbf{t}) f_{D,X}(\mathbf{x}, \mathbf{t}) dt \right) d\nu_{\mathcal{F}}(\mathbf{x}), \quad (4.9)$$

where $(\mathbf{x}, \mathbf{t}) = (x_1, \dots, x_k, t_1, \dots, t_k)$ are coordinates on V^{α} , $\{\xi_{\alpha}\}$ a partition of unity subordinate to the cover of \mathcal{S}^k by $\{V^{\alpha}\}$, and $d\mathbf{t}$ is induced from the usual Riemannian length measure along the orbits of X .

Remark 4.3 (Illustration for proof of Key Lemma). Let us illustrate our strategy in the case of the simplest diagram $D = \Theta$. For a fixed ϕ_X -invariant measure μ on \mathcal{S} , the question

is whether the following time average is μ -integrable:

$$\lambda_D(x) = \lim_{T \rightarrow \infty} \frac{1}{T^2} I_D(\mathcal{O}_T(x)), \quad x \in \mathcal{S}$$

For simplicity, here we disregard short paths.

Considering a finite cover of \mathcal{S}^2 by flowboxes V^α (defined for $k = 2$ in (4.8)), formula (4.9) tells us that it suffices to prove that $f_{D,X}$ is locally integrable with respect to $d\mathbf{t} \times \nu_{\mathcal{F}}$ in each V^α . Away from the diagonal $\Delta_{\{1,2\}}$ of \mathcal{S}^2 , $f_{D,X}$ is smooth, and so the hardest case is that of flowboxes intersecting $\Delta_{\{1,2\}}$. Without loss of generality consider a flowbox V^α , $\alpha = (\alpha_1, \alpha_2)$ with $\alpha_1 = \alpha_2$. To simplify the notation we denote it by $V = (\mathcal{T} \times I) \times (\mathcal{T} \times I)$, (where $\mathcal{T} \times I$ is a flowbox of X in \mathcal{S} , with $I = (-\epsilon, \epsilon)$). Let $F : \mathcal{T} \times \mathcal{T} \rightarrow \mathbb{R}$ be defined by

$$F(x, y) = \int_{I(x,y)} f_{D,X}(x, y, \mathbf{t}) d\mathbf{t}, \quad (4.10)$$

where

$$I(x, y) = \{x\} \times \{y\} \times I \times I \subset V, \quad I = (-\epsilon, \epsilon).$$

Since $C[2; \mathcal{S}] \subset C[2; \mathbb{R}^3]$ and $C[2; \mathbb{R}^3]$ is obtained by blowing up the diagonal $\Delta_{\{1,2\}}$ of $(\mathbb{R}^3)^2$, we can construct the metric \tilde{g} on $C[2; \mathbb{R}^3]$ from the standard Euclidean metric of \mathbb{R}^3 and pull it back to $C[2; \mathcal{T} \times I]$ via the map $\tilde{\phi}^2$ induced from the product flow $\phi^2 = \phi \times \phi$. The resulting metric on $C[2; \mathcal{T} \times I]$ will also be denoted by \tilde{g} . Recall that $\varpi_D = \varpi_{\Theta}$ is a smooth 2-form on $C[2; \mathcal{S}] \subset C[2; \mathbb{R}^3]$, defined via the Gauss map in (3.5). Thanks to Proposition 3.4, it pulls back to a smooth form on $C[2; \mathcal{T} \times I]$. The resulting pullback form will also be denoted by ϖ_D . In the case of configurations of two points, $C[2; \mathcal{T} \times I]$, the blowup map (3.1) can be set equal to the map defined in (2.2), namely

$$\beta : C(2; \mathcal{T} \times I) \longrightarrow C[2; \mathcal{T} \times I] = \text{Bl}(\mathcal{T} \times I, \Delta_{\{1,2\}})$$

Equations (4.2) and (4.3) imply

$$F(x, y) = \int_{I(x,y)} \alpha^* \varpi_D = \int_{\widetilde{I(x,y)}} \varpi_D, \quad (4.11)$$

where $\widetilde{I(x,y)}$ is the lift of $I(x, y) \subset V$ to $C[2; \mathcal{T} \times I]$. Using the metric \tilde{g} , Lemma 2.3 yields

$$|F(x, y)| \leq A_{\varpi_D, \tilde{g}} \text{vol}(\widetilde{I(x,y)}). \quad (4.12)$$

Claim: Volumes of lifts $\widetilde{I(x,y)}$ in the metric \tilde{g} are uniformly bounded over $\mathcal{T} \times \mathcal{T}$.

Given the claim, estimate (4.12) implies that F is pointwise bounded and thus $\nu_{\mathcal{F}}$ -integrable (because $\nu_{\mathcal{F}}$ is a finite measure). Applying this argument to each flow box chart $\{V^\alpha\}$, we can conclude that $f_{D,X}$ is $\bar{\mu}_\Delta$ -integrable as required.

Remark 4.4. One can regard the above reasoning as an alternative to the proof of Lemma 2.4 in [18, p. 1429].

Justification of Claim: The claim is intuitively clear, because the blowup map β “stretches” $I(x, y)$ locally by adding a “bump” (which is illustrated on the right side of Figure 5). To give a more precise argument, recall that $C[2; \mathcal{T} \times I]$ is a subspace of $C[2; \mathbb{R}^2 \times \mathbb{R}]$ and $C[2; \mathbb{R}^2 \times \mathbb{R}]$ is diffeomorphic to $(\mathbb{R}^2 \times \mathbb{R}) \times S^2 \times [0, \infty)$, (i.e. it is the complement of a tubular neighborhood of the thin diagonal). The blowup map

$$\beta : \left((\mathbb{R}^2 \times \mathbb{R})^2 - \Delta_{\{1,2\}} \right) \longrightarrow \left(C[2; \mathbb{R}^2 \times \mathbb{R}] - (\bar{\beta})^{-1}(\Delta_{\{1,2\}}) \right) \cong C(2; \mathbb{R}^2 \times \mathbb{R}),$$

can be written explicitly as

$$\beta : ((x, s), (y, t)) \longmapsto \left(\frac{x+y}{2}, \frac{s+t}{2}, \frac{(s-t, x-y)}{\sqrt{(s-t)^2 + |x-y|^2}}, \frac{1}{2} \sqrt{(s-t)^2 + |x-y|^2} \right).$$

Let $(x, y) \in \mathcal{T} \times \mathcal{T}$, $x \neq y$, and set $p = \frac{1}{2}(x+y)$, $q = \frac{1}{2}(x-y)$, conveniently changing variables to $u = \frac{1}{2}(s+t)$, $v = \frac{1}{2}(s-t)$, $s, t \in (-\epsilon, \epsilon)$. We obtain for $(u, v) \in (-\epsilon, \epsilon) \times (-\epsilon, \epsilon)$

$$((x, u), (y, v)) \longmapsto \left(p, u; \frac{(v, q)}{\sqrt{v^2 + |q|^2}}, \sqrt{v^2 + |q|^2} \right) =: (\psi_p(u); \psi_q(v)), \quad (4.13)$$

which, for a fixed x and y , gives a (u, v) -parametrization of the lift $\widetilde{I(x, y)}$. Here ψ_p, ψ_q denotes the curves given by respectively first and last two coordinates of the map (4.13). The volume $\text{vol}(\widetilde{I(x, y)})$ can now be estimated as

$$\text{vol}(\widetilde{I(x, y)}) \leq c_{\tilde{g}} \ell(\psi_p) (\ell(\psi_q) + 2\pi\epsilon),$$

where $\ell(\psi_p), \ell(\psi_q)$ are lengths of the curves ψ_p and ψ_q in the metric \tilde{g} and $c_{\tilde{g}}$ is a constant which depends only on \tilde{g} . Lengths $\ell(\psi_p)$ and $\ell(\psi_q)$ are proportional to ϵ and thus $(x, y) \longmapsto \text{vol}(\widetilde{I(x, y)})$ is uniformly bounded on $\mathcal{T} \times \mathcal{T}$ by a constant dependent only on the metric and the size of the flow box neighborhood. \square




FIGURE 5. Lift of orbits of a vector field on \mathbb{R}^2 (left) to $\text{Bl}(\mathbb{R}^2, \{x\}) \cong S^1 \times [1, +\infty)$ (right).

Proof of Key Lemma. Fix a flowbox chart $V^\alpha = \mathcal{T}^\alpha \times I^\alpha$ as defined in (4.8). It suffices to prove that the function

$$F_{D,\alpha}: \mathcal{T}^\alpha \longrightarrow \mathbb{R} \quad (4.14)$$

$$\mathbf{x} \longmapsto \int_{I^\alpha} f_{D,X}(\mathbf{x}, \mathbf{t}) d\mathbf{t}$$

is bounded for any α . Then, because the atlas $\{V^\alpha\}$ is finite, (4.9) implies $|\int_{\mathcal{S}^k} f_{D,X} \bar{\mu}_\Delta| \leq \infty$ as required. Note that $f_{D,X}$ is smooth away from the diagonals Δ_Q of \mathcal{S}^k , where $Q \subset \{1, \dots, k\}$, $\#Q \geq 2$, and

$$\Delta_Q = \{(x_1, \dots, x_k) \in \mathcal{S}^k \mid x_i = x_j \text{ for } i, j \in Q\}.$$

Let $\Delta = \bigcup_Q \Delta_Q$. We can cover the ϵ -neighborhood of Δ by open sets

$$V_Q^\alpha = \prod_{i=1}^k V_{\alpha_i}, \quad \text{where } \alpha_i = \alpha_j, \quad \text{for } i, j \in Q, \quad (4.15)$$

$$\text{and } \bar{V}_{\alpha_i} \cap \bar{V}_{\alpha_j} = \emptyset, \quad \text{for } i \in Q, j \notin Q.$$

Then the sup-norm of $f_{D,X}$ is bounded on the complement of the ϵ -neighborhood of Δ by some constant which only depends on X and ϖ_D ; see (4.2). Generally, we want to pick ϵ much smaller than ϵ , which is the size of the flow box charts. Thus, it suffices to prove that the functions $F = F_{D,\alpha}$ are bounded on V_Q^α for any α and Q . Up to a permutation of factors, suppose that $Q = \{1, \dots, r\} \subset \{1, \dots, k\}$, $2 \leq r \leq k$, $r = \#Q$ and $\alpha = (\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_{k-r})$ (for $\alpha = \alpha_1 = \dots = \alpha_r$). Then $V_Q^\alpha = \mathcal{T}^r \times I^r \times \mathcal{T}^\beta \times I^\beta$, where

$$\mathcal{T} = \mathcal{T}_\alpha, \quad I = I_\alpha, \quad \mathcal{T}^\beta = \mathcal{T}_{\beta_1} \times \dots \times \mathcal{T}_{\beta_{k-r}}, \quad \text{and } I^\beta = I_{\beta_1} \times \dots \times I_{\beta_{k-r}}.$$

Proposition 3.4 implies that the flow $\phi = \phi_X$ of X restricted to the flow box $\mathcal{T} \times I$ lifts to an embedding $\tilde{\phi}: C[r; \mathcal{T} \times I] \longrightarrow C[r; \mathbb{R}^3]$, which, by the second condition in (4.15), extends trivially to the embedding

$$\tilde{\phi}_Q: W_Q^\alpha \longrightarrow C[k; \mathbb{R}^3], \quad W_Q^\alpha := C[r; \mathcal{T} \times I] \times \mathcal{T}^\beta \times I^\beta.$$

Let $\bar{\alpha}_Q: W_Q^\alpha \longrightarrow V_Q^\alpha$ be the obvious projection induced by the restriction of the blowdown map (of Definition 3.3). Recall that for any $\mathbf{x} \in \mathcal{T}^r \times \mathcal{T}^\beta$, the lift $\widetilde{I_Q(\mathbf{x})}$ of $I_Q(\mathbf{x}) = \{\mathbf{x}\} \times I^r \times I^\beta$ to W_Q^α equals to the closure of $\bar{\alpha}_Q^{-1}(I_Q(\mathbf{x}))$ in W_Q^α . The k -form ϖ_D (3.9) pulls back to a smooth form on W_Q^α , and we may also pull back the metric \tilde{g} from $C[k; \mathbb{R}^3]$ to W_Q^α . By (4.2) and (4.14), a point value of F , for any $\mathbf{x} \in \mathcal{T}^r \times \mathcal{T}^\beta$, is given as

$$F(\mathbf{x}) = \int_{\widetilde{I_Q(\mathbf{x})}} \varpi_D. \quad (4.16)$$

Using Lemma 2.3, for some universal constant A_D we obtain a bound

$$|F(\mathbf{x})| \leq A_D \text{vol}(\widetilde{I_Q(\mathbf{x})}). \quad (4.17)$$

Therefore, analogously to what is outlined in Remark 4.3, it suffices to show that $\text{vol}(\widetilde{I_Q(\mathbf{x})})$ is uniformly bounded over $\mathcal{T}^r \times \mathcal{T}^\beta$. This is intuitively clear because $\text{vol}(I_Q(\mathbf{x}))$ is uniformly bounded by a constant proportional to ϵ^k (c.f. (4.8)) and $C[k; \mathbb{R}^3]$ is obtained from $(\mathbb{R}^3)^k$ by a sequence of blowups. Hence the philosophy presented in Remark 4.3 implies that $\text{vol}(\widetilde{I_Q(\mathbf{x})})$ is uniformly bounded as well. The remaining part of this proof provides details of this intuitive claim.

Summarizing, given a bounded ‘‘flow box’’: $W_Q^\alpha = C[r; \mathcal{T} \times I] \times \mathcal{T}^\beta \times I^\beta$ embedded via the flow in $C[k; \mathbb{R}^3]$, we intend to estimate the volume of the lift of $I_Q(\mathbf{x}) = \{\mathbf{x}\} \times I^r \times I^\beta \subset (\mathcal{T}^r \times \mathcal{T}^\beta) \times I^r \times I^\beta = V_Q^\alpha$ to W_Q^α for every $\mathbf{x} \in \mathcal{T}^r \times \mathcal{T}^\beta$. Specifically, we consider V_Q^α sitting in $\mathbb{R}^2 \times \mathbb{R}$ where the I factors of V_Q^α are mapped into the \mathbb{R} factors under the inclusion, and the corresponding blowup map

$$\begin{aligned} \alpha : C(k; \mathbb{R}^2 \times \mathbb{R}) &\longrightarrow A[k; \mathbb{R}^2 \times \mathbb{R}], \\ A[k; \mathbb{R}^2 \times \mathbb{R}] &= (\mathbb{R}^2 \times \mathbb{R})^k \times \prod_{S \subset \{1, \dots, k\}, |S| \geq 2} \text{Bl}((\mathbb{R}^2 \times \mathbb{R})^S, \Delta_S). \end{aligned} \quad (4.18)$$

Recall from Section 3 that $C[k; \mathbb{R}^2 \times \mathbb{R}]$ is obtained as a closure of the graph of the above map. The projection of the map α to the first factor of $A[k; \mathbb{R}^2 \times \mathbb{R}]$ is just the inclusion and the projections restricted to the $\text{Bl}(\cdot)$ factors are determined by the blowup maps as in (2.2). The metric on each $\text{Bl}_S = \text{Bl}((\mathbb{R}^2 \times \mathbb{R})^k, \Delta_S)$, further denoted by $\widetilde{g}_S(\varepsilon)$, is obtained via the construction of Section 2. The parameter ε is set to be sufficiently small, in particular smaller than the diameter of any flowbox chart. Recall that Bl_S is diffeomorphic to the complement of a tubular neighborhood of the thin diagonal Δ_S in $(\mathbb{R}^2 \times \mathbb{R})^S$, namely

$$\text{Bl}_S \cong (\mathbb{R}^2 \times \mathbb{R}) \times \mathbb{S}^{3|S|-4} \times [0, \infty). \quad (4.19)$$

and the map α restricted to factors of $A[k; \mathbb{R}^2 \times \mathbb{R}]$, indexed by $S = \{s_1, \dots, s_{|S|}\}$, can be specifically chosen as

$$\mathbf{y} = (y_1, \dots, y_{|S|}) \longmapsto \left(y_1; \frac{y_1 - y_2}{|\mathbf{y}'|}, \dots, \frac{y_1 - y_{|S|}}{|\mathbf{y}'|}; |\mathbf{y}'| \right), \quad \mathbf{y}' = (y_1 - y_2, \dots, y_1 - y_{|S|}). \quad (4.20)$$

This gives an embedding into the interior of Bl_S , i.e. into $(\mathbb{R}^2 \times \mathbb{R}) \times \mathbb{S}^{3|S|-4} \times (0, \infty)$.

For simplicity, suppose $\mathbf{x} \in \mathcal{T}^k \cong \mathcal{T}^r \times \mathcal{T}^\beta$ and $\mathbf{x} \in C(k; \mathcal{T})$, i.e. \mathbf{x} is away from the thick diagonal of \mathcal{T}^k . The restriction of the map α to $I(\mathbf{x}) = \{\mathbf{x}\} \times I^k$,¹⁰ gives a parametrization of the lift $\widetilde{I(\mathbf{x})}$ in $C[k; \mathbb{R}^2 \times \mathbb{R}]$. Let us denote this parametrization by $\gamma(t_1, \dots, t_k)$, where $\mathbf{t} = (t_1, \dots, t_k)$ are the variables of I^k and $\widetilde{I(\mathbf{x})} = \gamma(I(\mathbf{x}))$. Further let

$$X_i(\mathbf{t}) = \frac{\partial}{\partial t_i} \gamma(\mathbf{t}), \quad i = 1, \dots, k,$$

¹⁰Where we abbreviate $I^r \times I^\beta$ to I^k with $k = r + |\beta|$.

be images in $A[k; \mathbb{R}^2 \times \mathbb{R}]$ of coordinate vector fields under the derivative $D\gamma$. Then, for $\widetilde{I_Q}(\mathbf{x}) = \widetilde{\phi} \circ \alpha(I(\mathbf{x}))$ we have

$$\text{vol}(\widetilde{I_Q}(\mathbf{x})) \leq c_\phi \text{vol}(\widetilde{I}(\mathbf{x})) = c_\phi \int_{I^k} (|X_1 \wedge \cdots \wedge X_k|_{\widetilde{g}})^{\frac{1}{2}} dt, \quad (4.21)$$

where c_ϕ accounts for the C^1 -norm of the map $\widetilde{\phi}$. Each vector X_i has coordinates (X_i^j, X_i^S) , where $j = 1, \dots, k$ indexes factors: $(\mathbb{R}^2 \times \mathbb{R})^k$ in $A[k; \mathbb{R}^2 \times \mathbb{R}]$ and $S \subset \{1, \dots, k\}$, $|S| \geq 2$, indexes the Bl_S factors, we call j the *front index* and S the *set index* in the above decomposition of X_i . Substituting $X_i = \sum_{\mathbf{m}} X_i^{\mathbf{m}}$, where \mathbf{m} ranges over both j and S type indices, we estimate

$$\begin{aligned} |X_1 \wedge \cdots \wedge X_k|_{\widetilde{g}} &\leq \sum_{\mathbf{m}=(\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_k)} |X_1^{\mathbf{m}_1} \wedge \cdots \wedge X_k^{\mathbf{m}_k}|_{\widetilde{g}} \\ &\leq \sum_{\mathbf{m}} \prod_{l=1}^k |X_l^{\mathbf{m}_l}|_{\widetilde{g}} \leq \sum_{\mathbf{m}} \frac{1}{k} \sum_{l=1}^k (|X_l^{\mathbf{m}_l}|_{\widetilde{g}})^k, \end{aligned} \quad (4.22)$$

where the last step is a consequence of the arithmetic mean and Jensen's inequality (c.f. [32]). Consequently, estimating the integral in (4.21), boils down to estimating integrals in the form

$$\mathcal{J}(\mathbf{m}_l) = \int_{I^k} (|X_l^{\mathbf{m}_l}|_{\widetilde{g}})^k dt, \quad l = 1, \dots, k.$$

Without loss of generality (as we may always change the order of integration) suppose $l = 1$, and let

$$\mathcal{J} = \int_I \cdots \left(\int_I (|X_1^{\mathbf{m}_1}(t_1, \dots, t_k)|_{\widetilde{g}})^k dt_1 \right) \cdots dt_k.$$

For a fixed $\mathbf{t}_0 = (t_2, \dots, t_k)$, the inner integral:

$$E_k(\gamma_{\mathbf{m}_1}) = \int_I (|X_1^{\mathbf{m}_1}(t_1, \mathbf{t}_0)|_{\widetilde{g}})^k dt_1,$$

represents the L^k -energy¹¹ of the curve parametrized by $\gamma_{\mathbf{m}_1} : t_1 \rightarrow \pi_{\mathbf{m}_1}(\gamma(t_1, \mathbf{t}_0))$, where $\pi_{\mathbf{m}_1}$ is a projection onto the \mathbf{m}_1 -coordinate of $A[k; \mathbb{R}^2 \times \mathbb{R}]$. If \mathbf{m}_1 is a front index then $|X_1^{\mathbf{m}_1}(t_1, \mathbf{t}_0)|_{\widetilde{g}} = 1$ and $E_k(\gamma_{\mathbf{m}_1}) \leq \ell(\gamma_{\mathbf{m}_1})$. Since $\ell(\gamma_{\mathbf{m}_1}) \leq c_\epsilon \epsilon$, for some constant $c_\epsilon > 0$, we obtain

$$\mathcal{J}(\mathbf{m}_1) \leq (c_\epsilon \epsilon)^k.$$

In the case $\mathbf{m}_1 = S$ is a set index, the map $\gamma_{\mathbf{m}_1}$ parametrizes a curve in Bl_S , which projected via (4.20) onto the $\mathbb{S}^{3|S|-4}$ factor is a "piece" of a great circle. Then a simple computation in the metric $\widetilde{g}_S(\varepsilon)$ leads to the following estimate

$$E_k(\gamma_{\mathbf{m}_1}) \leq c_{k,\epsilon} (2\pi + 1)^k \epsilon^k,$$

where we used $\varepsilon \ll \epsilon$, again $\mathcal{J}(\mathbf{m}_1)$ is uniformly bounded. Applying $\int_{I^k} (\cdot)$ to both sides of (4.22) we obtain from (4.21) that $\text{vol}(\widetilde{I_Q}(\mathbf{x}))$ is estimated by a sum of \mathcal{J} -type terms. Therefore

¹¹i.e. the L^k -norm to the k th power.

using estimates for $J(\mathbf{m}_1)$ we obtain the required uniform bound

$$\text{vol}(\widetilde{I_Q(\mathbf{x})}) \leq c_{k,\epsilon,\phi}(1 + 2\pi\epsilon)^k(\epsilon)^{2k}.$$

In the case \mathbf{x} belongs to the thick diagonal of \mathcal{T}^k , we obtain the above bound by considering \mathbf{x} as a limit of points from $C(k; \mathcal{T})$. \square

4.2. Short paths. We are now ready to show that the short paths do not contribute to the limits in (4.1) and (4.4).

Lemma 4.5. *We have*

$$\bar{\lambda}_D(x) = \lambda_D(x), \quad a.e. \quad (4.23)$$

Proof. We will adapt the classical argument of Arnold from [4].¹² Recall that the short curves are denoted by $\{\sigma(x, y)\}_{x,y \in \mathcal{S}}$. The difference of integrals in the limits defining λ_D (4.5) and $\bar{\lambda}_D$ (4.1) respectively is a sum of the following terms (up to permutation of \int_0^T and \int_0^1) for $i \geq 1$, and $i + j = k$:

$$J_{i,j} = \underbrace{\int_0^1 \cdots \int_0^1}_i \underbrace{\int_0^T \cdots \int_0^T}_j f_{D,X}(\sigma_{x,T}(s_1), \cdots, \sigma_{x,T}(s_i), \phi(x, t_1) \cdots, \phi(x, t_j)) dt ds,$$

where $\sigma_{x,T} : [0, 1] \rightarrow \mathcal{S}$ is a parametrization of $\sigma(x, \phi(x, T))$. Here \int_0^T is an integral over the orbit of X and \int_0^1 is an integral over the short path segment, from (4.4). Fixing small enough $\epsilon > 0$, we may subdivide each $[0, T]$ so that the integral \int_0^T is roughly the sum $\int_0^\epsilon + \int_\epsilon^{2\epsilon} + \cdots + \int_{\epsilon(\lceil \frac{1}{\epsilon} T \rceil - 1)}^{\epsilon \lceil \frac{1}{\epsilon} T \rceil}$, and each ϵ -interval $[\epsilon(k-1), \epsilon k]$, $1 \leq k \leq \lceil \frac{T}{\epsilon} \rceil$, parametrizes a piece of an orbit within a flowbox chart of X . Analogously, we may subdivide the unit intervals parametrizing the short paths and therefore split the \int_0^1 integral into the ϵ -pieces, also fitting into flowbox neighborhoods of X . Let the index k_l , $1 \leq l \leq j$, enumerate the sums for the \int_0^T 's and the index m_z , $1 \leq z \leq i$, enumerate the sums for \int_0^1 's. Then, the above formula for $J_{i,j}$ yields

$$|J_{i,j}| \leq \sum_{\substack{m_1, \dots, m_i \\ k_1, \dots, k_j}} \left| \int_{\epsilon(m_1-1)}^{\epsilon m_1} \cdots \int_{\epsilon(m_i-1)}^{\epsilon m_i} \int_{\epsilon(k_1-1)}^{\epsilon k_1} \cdots \int_{\epsilon(k_j-1)}^{\epsilon k_j} f_{D,X} dt ds \right|.$$

Each integral term in the above sum can be expressed, similarly as in (4.16), as an integral over a lift of the product of the short ϵ -pieces of σ 's and the orbits of X , over a smooth differential form ϖ_D on $C[k; \mathbb{R}^3]$. Therefore, by the estimate (4.17) each integral in the above sum can be bounded above by a constant $A_{X,D}$ which only depends on the vector field, ϖ_D , and the metric. Since the number of terms in the sum $J_{i,j}$ is given by $(\lceil 1/\epsilon \rceil)^i (\lceil T/\epsilon \rceil)^j$ we obtain

$$|J_{i,j}| \leq A_{X,D} (\lceil 1/\epsilon \rceil)^i (\lceil T/\epsilon \rceil)^j.$$

¹²One needs to be cautious about this argument in the case the domain of the vector field cannot be covered by finitely many flowbox type neighborhoods; see Example 4 in [43].

Since there are $\binom{k}{i}$ terms of type $J_{i,j}$ in the difference $\bar{\lambda}_D - \lambda_D$ and $j = k - i$, we obtain, for any $x \in \mathcal{S}$,

$$|\bar{\lambda}_D(x) - \lambda_D(x)| \leq A_{X,D} \lim_{T \rightarrow \infty} \frac{1}{T^k} \sum_{i=1}^k \binom{k}{i} ([1/\epsilon])^i ([T/\epsilon])^{k-i} = 0. \quad \square$$

4.3. Proofs of Theorems A, Corollary A, and Theorem B. Recall the statements of these three results from the Introduction.

Proof of Theorem A and Corollary A. Part (i) has already been proven in Key Lemma. For part (ii), following Theorem 3.6, any finite type n invariant V_W is a linear combination of integrals I_D of (3.7). Specifically, for appropriate coefficients $a_D \in \mathbb{R}$ and $D_1 = \Theta$, we can express V_W as

$$V_W(K) = \sum_{k=1}^{2n} J_k(K) + b I_{D_1}(K), \quad \text{for } J_k(K) = \sum_{D \in TD_n; k(D)=k} a_D I_D(K), \quad K \in \mathcal{K}. \quad (4.24)$$

In order to observe the almost everywhere convergence in

$$\bar{\lambda}_W(x) = \lim_{T \rightarrow \infty} \frac{1}{T^{2n}} V_W(\bar{\mathcal{O}}_T(x)),$$

we take the corresponding linear combination of T^{2n} -time averages of terms in (4.24). Namely, we have

$$\bar{\lambda}_W(x) = \sum_{k=1}^{2n} \lim_{T \rightarrow \infty} \frac{1}{T^{2n}} J_k(\bar{\mathcal{O}}_T(x)) + b \lim_{T \rightarrow \infty} \frac{1}{T^{2n}} I_{D_1}(\bar{\mathcal{O}}_T(x)). \quad (4.25)$$

By Key Lemma, for $n > 1$, the terms in the sum (4.25) with $k < n$ vanish in the limit as does the I_{D_1} term. As a result, we have

$$\bar{\lambda}_W(x) = \bar{\lambda}_{W^{2n}}(x) = \sum_{D \in TD_n; k(D)=2n} a_D \bar{\lambda}_D(x).$$

Further, if $J_{2n}(\bar{\mathcal{O}}_T(x))$ is $o(T^{2n-1})$, then we may consider T^{2n-1} -time averages of V_W and obtain

$$\bar{\lambda}_W(x) = \bar{\lambda}_{W^{2n-1}}(x) = \sum_{D \in TD_n; k(D)=2n-1} a_D \bar{\lambda}_D(x).$$

This reasoning further applies, if the terms J_k are of lower order, and this therefore gives the proof of Corollary A.

It remains to prove invariance under volume-preserving deformations as claimed in (iii). Note that, given $h \in \text{Diff}_0(\mathbb{R}^3, \mu)$, the short path system $h\Sigma = \{h \circ \sigma\}$ on $\mathcal{S}' = h(\mathcal{S})$ obtained from $\Sigma = \{\sigma\}$ has the same properties as the original system Σ on \mathcal{S} with respect to the pulled-back metric on \mathcal{S}' . In particular, Lemma 4.5 holds for $h\Sigma$. Now, for any $T > 0$ and $x \in \mathcal{S}$, consider knots $K_{h_*X} = \bar{\mathcal{O}}_T^{h_*X}(x)$ and $K_X = \bar{\mathcal{O}}_T^X(h^{-1}(x))$ (where we used $h\Sigma$ to close

up K_{h_*X} and Σ to close up K_X). By (1.6), we have

$$K_{h_*X} = h(K_X).$$

Since $h \in \text{Diff}_0(\mathbb{R}^3, \mu)$, we conclude that K_{h_*X} and K_X are isotopic, implying

$$V_W(K_{h_*X}) = V_W(K_X).$$

Taking $\lim_{T \rightarrow \infty} \frac{1}{T^{2n}}(\cdot)$ of both sides in the above equation yields

$$\bar{\lambda}_W^{h_*X}(x) = \bar{\lambda}_W^X(h^{-1}(x)), \quad a.e.$$

After a change of variables (using the fact that h is μ -preserving), we obtain

$$\mathcal{V}_W(h_*X) = \mathcal{V}_W(X). \quad \square$$

From the above argument, observe that $\lambda_{W^k}(x)$ is a time average of the L^1 -functions

$$f_{W,X,k} : C(2n; \mathbb{R}^3) \longrightarrow \mathbb{R}, \quad f_{W,X,k} := \sum_{D \in TD_n; k(D)=k} a_D f_{D,X,k},$$

for $f_{D,X,k} = f_{D,X}$ as defined in (4.2). Applying the Multiparameter Ergodic Theorem [11, 38] to $f_{W,X,k}$, we obtain the following formula for the vector field invariant $\mathcal{V}_{W,k} : \text{Vect}(\mathcal{S}, \mu) \longrightarrow \mathbb{R}$:

$$\mathcal{V}_{W,k}(X) = \int_{\mathcal{S}^k} f_{W,X,k} \bar{\mu}_\Delta \tag{4.26}$$

(recall $\bar{\mu}_\Delta$ is a diagonal invariant measure on \mathcal{S}^k given in (4.6)).




FIGURE 6. A top degree diagram perturbation leads to pairwise linking number diagrams.

Proof of Theorem B. By assumption, the domain \mathcal{S} is equipped with the standard volume form μ and X is an ergodic μ -preserving nonvanishing vector field. For simplicity, we assume that μ induces a probability measure on \mathcal{S} . Ergodicity of X on \mathcal{S} implies, among other things, that almost every orbit of X densely fills the interior of \mathcal{S} . Clearly, μ induces a ϕ_X^k -invariant ergodic probability measure on \mathcal{S}^k via the $3k$ -form $\mu^k = \underbrace{\mu \times \cdots \times \mu}_{k \text{ times}}$. By Key Lemma, $f_{W,X}$ is in $L^1(\mu^{2n})$, and thus the ergodicity of the ϕ_X^{2n} -action implies that the integral

$$\int_{\mathcal{S}^{2n}} f_{W,X} \mu^{2n}$$

equals

$$\lim_{T \rightarrow \infty} \frac{1}{T^{2n}} \underbrace{\int_0^T \cdots \int_0^T}_{2n \text{ times}} f_{W^{2n}, X}(\phi(x_1, t_1), \dots, \phi(x_{2n}, t_{2n})) dt_1 \cdots dt_{2n}, \quad (4.27)$$

for almost every point $\mathbf{x} = (x_1, \dots, x_{2n}) \in \mathcal{S}^{2n}$. Choosing \mathbf{x} to be away from the thick diagonal, we have $2n$ distinct orbits $\bar{\mathcal{O}}_T(\mathbf{x}) = \bar{\mathcal{O}}_T(x_1) \times \cdots \times \bar{\mathcal{O}}_T(x_{2n})$ through each coordinate point. For each top degree diagram (i.e. a chord diagram) $D \in \mathcal{D}_n$, $k(D) = 2n$, the integral of the associated differential form ϖ_D over $\bar{\mathcal{O}}_T(\mathbf{x})$ splits as a product of linking numbers of pairs of points associated with the chords of D . This can be thought of as a perturbation of the diagram, as the vertices are no longer on the same orbit; see Figure 6 for an illustration. Explicitly, for $\bar{\mathcal{O}}_T(\mathbf{x})$ and $\varpi_D = \prod_{(i,j) \in \mathcal{E}(D)} \omega_{i,j}$, from (4.2) and the fact that $\int_{\bar{\mathcal{O}}_T(x_i) \times \bar{\mathcal{O}}_T(x_j)} \omega_{i,j} = \text{lk}(\bar{\mathcal{O}}_T(x_i), \bar{\mathcal{O}}_T(x_j))$, we have (up to short paths)

$$\underbrace{\int_0^T \cdots \int_0^T}_{2n \text{ times}} f_{D, X}(\phi(x_1, t_1), \dots, \phi(x_{2n}, t_{2n})) dt_1 \cdots dt_{2n} = \prod_{(i,j) \in \mathcal{E}(D)} \text{lk}(\bar{\mathcal{O}}_T(x_i), \bar{\mathcal{O}}_T(x_j)). \quad (4.28)$$

By definition of $\mathcal{H}(X)$ (see (1.3)) and the ergodicity assumption, summing up over all top order diagrams $D \in \mathcal{D}_n$, we obtain from (4.27) the independence of the limit of short paths and from (4.28) we obtain

$$\int_{\mathcal{S}^{2n}} f_{W, X} \mu^{2n} = c_W (\mathcal{H}(X))^n, \quad (4.29)$$

where c_W is a constant independent of X .

Next, we turn to the proof of the identity in (1.12). Observe that in the space of probability measures $\mathcal{M}(\mathcal{S}^{2n})$, the diagonal measure μ_Δ can be approximated by a sequence of probability measures supported on the δ -tubular neighborhood $U_\delta = U_\delta(\Delta)$ of the thin diagonal Δ of \mathcal{S}^{2n} . These measures can be precisely defined as

$$\nu_\delta^{2n} = \frac{\chi_\delta}{\text{vol}(U_\delta)} \nu^{2n}, \quad \chi_\delta(\mathbf{x}) = \begin{cases} 1, & \mathbf{x} \in U_\delta, \\ 0, & \mathbf{x} \notin U_\delta. \end{cases}$$

Since $\mu_\delta^{2n} \rightarrow \mu_\Delta$, $\delta \rightarrow 0$ in $\mathcal{M}(\mathcal{S}^{2n})$. Thanks to the weak compactness of $\mathcal{M}(\mathcal{S}^{2n})$, the sequence of the associated invariant measures $\bar{\mu}_\delta^{2n}$, built via the formula (4.6), converges to the diagonal invariant measure $\bar{\mu}_\Delta$ in $\mathcal{M}(\mathcal{S}^{2n})$. From Key Lemma, for each δ , $f_{W, X}$ is in $L^1(\bar{\mu}_\delta^{2n})$. Since the right hand side in (4.27) is independent of the choice of \mathbf{x} (as long as it is generic), for a given δ we may suppose $\mathbf{x} \in U_\delta$ and obtain from (4.29) and the assumption of ergodicity the identity

$$\int_{\mathcal{S}^{2n}} f_{W, X} \bar{\mu}_\delta^{2n} = \lim_{T \rightarrow \infty} \frac{1}{T^{2n}} \underbrace{\int_0^T \cdots \int_0^T}_{2n \text{ times}} f_{W, X}(\mathbf{x}, \mathbf{t}) dt = c_W (\mathcal{H}(X))^n.$$

Since $\bar{\mu}_\delta^{2n} \rightarrow \bar{\mu}_\Delta$ in $\mathcal{M}(\mathcal{S}^{2n})$, we deduce (1.12). The second part of Theorem B can be justified analogously. \square

5. QUADRATIC HELICITY, ENERGY, AND PROOF OF THEOREM C.

The methods presented in the previous sections can be applied almost without any changes to the setting of asymptotic links. One difference between the case of knots and links is a choice of the diagonal invariant measure $\bar{\mu}_\Delta$ in (4.6). Rather than presenting this obvious generalization, the rest of this section is devoted to an illustration of the relevant constructions for the simplest finite type 2 invariant associated with a 2–component link, the square of the linking number lk^2 . We observe that in the setting of asymptotic links, lk^2 leads to *quadratic helicity* that was recently proposed by Akhmetiev in [1]. Further, it is the simplest invariant that can provide a sharper lower bound for the fluid energy than $\mathcal{H}(X)$, as claimed in Theorem C.

The weight system associated to lk^2 is given by just one trivalent diagram which we denote by D_{lk^2} , pictured in Figure 7. The configuration of points and chords on D_{lk^2} implies




FIGURE 7. A trivalent diagram D_{lk^2} for lk^2 .

a choice of the invariant measure on \mathcal{S}^4 associated with the flow of X . Namely, we start with the product $\phi_X^2 = \phi_X \times \phi_X$ –invariant measure $\mu \times \mu$ on $\mathcal{S} \times \mathcal{S}$ and push it forward to the 4–fold product \mathcal{S}^4 by the inclusion $j: (x, y) \mapsto (x, x, y, y)$. Let us denote the diagonal, parametrized by j , by $\Delta_{(2)} = \Delta_{\{\{1,3\},\{2,4\}\}}$. Also denote the pushforward measure by $\mu_{\Delta_{(2)}}$ and the associated ϕ_X^4 –invariant measure by $\bar{\mu}_{\Delta_{(2)}}$ (i.e. $\bar{\mu}_{\Delta_{(2)}} = \int \mu_{\Delta_{(2)}}$). By virtue of Theorem A, the asymptotic invariant of X associated with lk^2 equals the quadratic helicity of [1] and is by (4.26) given as

$$\mathcal{H}^2(X) = \int_{\mathcal{S}^4} \varpi_{D_{\text{lk}^2}}(X, X, X, X) \bar{\mu}_{\Delta_{(2)}}, \tag{5.1}$$

where

$$\varpi_{D_{\text{lk}^2}} = \alpha^* \omega_{1,2} \wedge \alpha^* \omega_{3,4}.$$

(because D_{lk^2} has no free vertices). Observe that $\mathcal{H}^2(X) \geq 0$, whereas $\mathcal{H}(X)$ can be negative. We can easily show examples when $\mathcal{H}(X) = 0$ but $\mathcal{H}^2(X) > 0$ (see [4, p. 344]). Therefore it is of general interest to derive an analog of inequality (1.15) for $\mathcal{H}^2(X)$.

Proof of Theorem C. Recall [15] that the diagonal invariant measure $\bar{\mu}_{\Delta(2)}$ can be arbitrarily well approximated in $\mathcal{M}(\mathbb{S}^4)$ by positive finite linear combinations

$$\bar{\mu}_n = \sum_{i=1}^n a_i \bar{\mu}_{\mathbf{x}_i}, \quad a_i > 0, \quad (5.2)$$

where $\bar{\mu}_{\mathbf{x}_i}$ is a ϕ_X^4 -invariant measure obtained from averaging a Dirac delta $\delta_{\mathbf{x}_i}$ supported at a point $\mathbf{x}_i = (x_i, x_i, y_i, y_i)$ on the diagonal $\Delta(2)$. More precisely, if $\mu_n = \sum_{i=1}^n a_i \delta_{\mathbf{x}_i}$ as an approximation of $\mu_{\Delta(2)}$, $\bar{\mu}_n = \bar{\int} \mu_n$ is an approximation of $\bar{\mu}_{\Delta(2)}$. In fact, approximating $\mu \times \mu$ by $\sum_{i=1}^n b_i \delta_{(x_i, y_i)}$, $b_i > 0$, and applying the pushforward under j we conclude that the coefficients in (5.2) are given as

$$a_i = b_i^2.$$

Note that each $\bar{\mu}_{\mathbf{x}_i}$ is a product measure, i.e.

$$\bar{\mu}_{\mathbf{x}_i} = \bar{\mu}_{(x_i, y_i)}^{\{1,2\}} \times \bar{\mu}_{(x_i, y_i)}^{\{3,4\}}, \quad (5.3)$$

where $\bar{\mu}_{(x_i, y_i)}^{\{k,l\}}$ is a pushforward of $\bar{\mu}_{(x_i, y_i)} = \bar{\int} \delta_{(x_i, y_i)}$ under the inclusion of $\mathbb{S} \times \mathbb{S}$ into the (k, l) -coordinates factor of $(\mathbb{S} \times \mathbb{S})^2 = \mathbb{S}^4$. By the proof of Theorem A, the function $f_W = \alpha^* \omega_{1,2} \wedge \alpha^* \omega_{3,4}(X^{\wedge 4})$ is $\bar{\mu}_n$ -integrable for each n . Moreover, if we set

$$f_{1,2} = \alpha^* \omega_{1,2}(X, X), \quad f_{3,4} = \alpha^* \omega_{3,4}(X, X),$$

then

$$f_W = \alpha^* \omega_{1,2} \wedge \alpha^* \omega_{3,4}(X^{\wedge 4}) = \alpha^* \omega_{1,2}(X, X) \alpha^* \omega_{3,4}(X, X) = f_{1,2} f_{3,4}.$$

Note that the functions $f_{1,2}$ and $f_{3,4}$ are constant on appropriate \mathbb{S}^2 factors of \mathbb{S}^4 . Using (5.2) and (5.3), we obtain

$$\begin{aligned} \left| \int_{\mathbb{S}^4} f_W \bar{\mu}_n \right| &= \left| \sum_{i=1}^n b_i^2 \left(\int_{\mathbb{S}^2} f_{1,2} \bar{\mu}_{(x_i, y_i)}^{\{1,2\}} \right) \left(\int_{\mathbb{S}^2} f_{3,4} \bar{\mu}_{(x_i, y_i)}^{\{3,4\}} \right) \right| \\ &\leq \left(\sum_{i=1}^n b_i \int_{\mathbb{S}^2} |f_{1,2}| \bar{\mu}_{(x_i, y_i)}^{\{1,2\}} \right) \left(\sum_{i=1}^n b_i \int_{\mathbb{S}^2} |f_{3,4}| \bar{\mu}_{(x_i, y_i)}^{\{3,4\}} \right). \end{aligned}$$

Passing to the limit in $\mathcal{M}(\mathbb{S}^4)$ as $n \rightarrow \infty$, we have $\mu_n \rightarrow \bar{\mu}_{\Delta(2)}$ and $\sum_i b_i \bar{\mu}_{(x_i, y_i)}^{\{k,l\}} \rightarrow \mu \times \mu$. Therefore

$$\mathcal{H}^2(X) \leq \left(\int_{\mathbb{S}^2} |\alpha^* \omega_{1,2}(X, X)| \mu \times \mu \right) \left(\int_{\mathbb{S}^2} |\alpha^* \omega_{3,4}(X, X)| \mu \times \mu \right) = c(X)^2, \quad (5.4)$$

where $c(X)$ stands for the asymptotic crossing number as defined in [22, p. 191], and the last identity is a consequence of $c(X) = \int_{\mathbb{S}^2} |\alpha^* \omega(X, X)| \mu \times \mu$ given in [22]. The estimate [22, Equation (1.9)]

$$E_{3/2}(X) \geq \left(\frac{16}{\pi} \right)^{1/4} c(X)^{3/4}$$

immediately yields the required bound in (1.17). \square

REFERENCES

- [1] P. M. Akhmetev. Quadratic helicities and the energy of magnetic fields. *Proceedings of the Steklov Institute of Mathematics*, 278(1):10–21, 2012.
- [2] P. M. Akhmetev. On a new integral formula for an invariant of 3-component oriented links. *J. Geom. Phys.*, 53(2):180–196, 2005.
- [3] D. Altschüler and L. Freidel. On universal Vassiliev invariants. *Comm. Math. Phys.*, 170(1):41–62, 1995.
- [4] V. I. Arnold. The asymptotic Hopf invariant and its applications. *Selecta Math. Soviet.*, 5(4):327–345, 1986. Selected translations.
- [5] V. I. Arnold. *Arnold’s problems*. Springer-Verlag, Berlin, 2004. Translated and revised edition of the 2000 Russian original, With a preface by V. Philippov, A. Yakivchik and M. Peters.
- [6] V. I. Arnold and B. A. Khesin. *Topological methods in hydrodynamics*, volume 125 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1998.
- [7] S. Baader. Asymptotic Rasmussen invariant. *C. R. Math. Acad. Sci. Paris*, 345(4):225–228, 2007.
- [8] S. Baader. Asymptotic concordance invariants for ergodic vector fields. *Comment. Math. Helv.*, 86(1):1–12, 2011.
- [9] S. Baader and J. Marché. Asymptotic Vassiliev invariants for vector fields. *Bulletin de la Société Mathématique de France*, 140(4):569–582, 2012.
- [10] D. Bar-Natan. On the Vassiliev knot invariants. *Topology*, 34(2):423–472, 1995.
- [11] M. E. Becker. Multiparameter groups of measure-preserving transformations: a simple proof of Wiener’s ergodic theorem. *Ann. Probab.*, 9(3):504–509, 1981.
- [12] P. Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley and Sons Inc., New York, second edition, 1999. A Wiley-Interscience Publication.
- [13] R. Bott and C. Taubes. On the self-linking of knots. *J. Math. Phys.*, 35(10):5247–5287, 1994. Topology and physics.
- [14] G. Călugăreanu. L’intégrale de Gauss et l’analyse des nœuds tridimensionnels. *Rev. Math. Pures Appl.*, 4:5–20, 1959.
- [15] A. Candel and L. Conlon. *Foliations. I*, volume 23 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2000.
- [16] J. Cantarella and J. Parsley. A new cohomological formula for helicity in \mathbb{R}^{2k+1} reveals the effect of a diffeomorphism on helicity. *J. Geom. Phys.*, 60(9):1127–1155, 2010.
- [17] S. Chmutov, S. Duzhin, and J. Mostovoy. *Introduction to Vassiliev knot invariants*. Cambridge University Press, Cambridge, 2012.
- [18] G. Contreras and R. Iturriaga. Average linking numbers. *Ergodic Theory Dynam. Systems*, 19(6):1425–1435, 1999.
- [19] N. W. Evans and M. A. Berger. A hierarchy of linking integrals. In *Topological aspects of the dynamics of fluids and plasmas (Santa Barbara, CA, 1991)*, volume 218 of *NATO Adv. Sci. Inst. Ser. E Appl. Sci.*, pages 237–248. Kluwer Acad. Publ., Dordrecht, 1992.
- [20] E. R. Fadell and S. Y. Husseini. *Geometry and topology of configuration spaces*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2001.
- [21] M. H. Freedman. Zeldovich’s neutron star and the prediction of magnetic froth. In *The Arnoldfest (Toronto, ON, 1997)*, volume 24 of *Fields Inst. Commun.*, pages 165–172. Amer. Math. Soc., Providence, RI, 1999.
- [22] M. H. Freedman and Z.-X. He. Divergence-free fields: energy and asymptotic crossing number. *Ann. of Math. (2)*, 134(1):189–229, 1991.

- [23] W. Fulton and R. MacPherson. A compactification of configuration spaces. *Ann. of Math. (2)*, 139(1):183–225, 1994.
- [24] J. Gambaudo and É. Ghys. Enlacements asymptotiques. *Topology*, 36(6):1355–1379, 1997.
- [25] J. M. Gambaudo and É. Ghys. Signature asymptotique d’un champ de vecteurs en dimension 3. *Duke Math. J.*, 106(1):41–79, 2001.
- [26] B. A. Khesin. Geometry of higher helicities. *Mosc. Math. J.*, 3(3):989–1011, 1200, 2003. {Dedicated to Vladimir Igorevich Arnold on the occasion of his 65th birthday}.
- [27] R. Komendarczyk. The third order helicity of magnetic fields via link maps. *Comm. Math. Phys.*, 292(2):431–456, 2009.
- [28] R. Komendarczyk. The third order helicity of magnetic fields via link maps. II. *J. Math. Phys.*, 51(12):122702, 16, 2010.
- [29] M. Kontsevich. Vassiliev’s knot invariants. In *I. M. Gelfand Seminar*, volume 16 of *Adv. Soviet Math.*, pages 137–150. Amer. Math. Soc., Providence, RI, 1993.
- [30] D. Kotschick and T. Vogel. Linking numbers of measured foliations. *Ergodic Theory Dynam. Systems*, 23(2):541–558, 2003.
- [31] P. Laurence and E. Stredulinsky. Asymptotic Massey products, induced currents and Borromean torus links. *J. Math. Phys.*, 41(5):3170–3191, 2000.
- [32] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [33] R. B. Melrose. *Differential Analysis on Manifolds with Corners*. –online notes available at <http://www-math.mit.edu/~rbm/book.html>.
- [34] H. K. Moffatt. The degree of knottedness of tangled vortex lines. *J. Fluid Mech*, 35(1):117–129, 1969.
- [35] T. Rivière. High-dimensional helicities and rigidity of linked foliations. *Asian J. Math.*, 6(3):505–533, 2002.
- [36] D. Ruelle and D. Sullivan. Currents, flows and diffeomorphisms. *Topology*, 14(4):319–327, 1975.
- [37] D. P. Sinha. Manifold-theoretic compactifications of configuration spaces. *Selecta Math. (N.S.)*, 10(3):391–428, 2004.
- [38] A. A. Tempelman. Ergodic theorems for general dynamical systems. *Dokl. Akad. Nauk SSSR*, 176:790–793, 1967.
- [39] D. P. Thurston. Integral Expressions for the Vassiliev Knot Invariants. *Harvard University senior thesis, April 1995*, arXiv math/9901110, 1999.
- [40] H. v. Bodecker and G. Hornig. Link invariants of electromagnetic fields. *Phys. Rev. Lett.*, 92(3):030406, 4, 2004.
- [41] V. A. Vassiliev. *Complements of discriminants of smooth maps: topology and applications*, volume 98 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1992. Translated from the Russian by B. Goldfarb.
- [42] A. Verjovsky and R. F. Vila Freyer. The Jones-Witten invariant for flows on a 3-dimensional manifold. *Comm. Math. Phys.*, 163(1):73–88, 1994.
- [43] T. Vogel. On the asymptotic linking number. *Proc. Amer. Math. Soc.*, 131(7):2289–2297 (electronic), 2003.
- [44] I. Volić. A survey of Bott-Taubes integration. *J. Knot Theory Ramifications*, 16(1):1–42, 2007.
- [45] L. Woltjer. On hydromagnetic equilibrium. *Proc. Nat. Acad. Sci. U.S.A.*, 44:833–841, 1958.

TULANE UNIVERSITY, NEW ORLEANS, LOUISIANA 70118

E-mail address: `rako@tulane.edu`

WELLESLEY COLLEGE, WELLESLEY, MA 02481-8203

E-mail address: `ivollic@wellesley.edu`