



$$\begin{aligned}
 2. (a) (i) L_6 &= \sum_{i=1}^6 f(x_{i-1}) \Delta x \quad [\Delta x = \frac{12-0}{6} = 2] \\
 &= 2[f(x_0) + f(x_1) + f(x_2) + f(x_3) + f(x_4) + f(x_5)] \\
 &= 2[f(0) + f(2) + f(4) + f(6) + f(8) + f(10)] \\
 &\approx 2(9 + 8.8 + 8.2 + 7.3 + 5.9 + 4.1) \\
 &= 2(43.3) = 86.6
 \end{aligned}$$




$$\begin{aligned}
 (ii) R_6 &= L_6 + 2 \cdot f(12) - 2 \cdot f(0) \\
 &\approx 86.6 + 2(1) - 2(9) = 70.6
 \end{aligned}$$

[Add area of rightmost lower rectangle
and subtract area of leftmost upper rectangle.]



$$\begin{aligned}
 (iii) M_6 &= \sum_{i=1}^6 f(x_i^*) \Delta x \\
 &= 2[f(1) + f(3) + f(5) + f(7) + f(9) + f(11)] \\
 &\approx 2(8.9 + 8.5 + 7.8 + 6.6 + 5.1 + 2.8) \\
 &= 2(39.7) = 79.4
 \end{aligned}$$



(b) Since f is decreasing, we obtain an *overestimate* by using *left* endpoints; that is, L_6 .

(c) Since f is decreasing, we obtain an *underestimate* by using *right* endpoints; that is, R_6 .

(d) M_6 gives the best estimate, since the area of each rectangle appears to be closer to the true area than the overestimates and underestimates in L_6 and R_6 .

$$13. \text{ Lower estimate for oil leakage: } R_5 = (7.6 + 6.8 + 6.2 + 5.7 + 5.3)(2) = (31.6)(2) = 63.2 \text{ L.}$$

$$\text{Upper estimate for oil leakage: } L_5 = (8.7 + 7.6 + 6.8 + 6.2 + 5.7)(2) = (35)(2) = 70 \text{ L.}$$

14. We can find an upper estimate by using the final velocity for each time interval. Thus, the distance d traveled after 62 seconds can be approximated by

$$d = \sum_{i=1}^6 v(t_i) \Delta t_i = (185 \text{ ft/s})(10 \text{ s}) + 319 \cdot 5 + 447 \cdot 5 + 742 \cdot 12 + 1325 \cdot 27 + 1445 \cdot 3 = 54,694 \text{ ft}$$

16. For an increasing function, using left endpoints gives us an underestimate and using right endpoints results in an overestimate.

We will use M_6 to get an estimate. $\Delta t = \frac{30-0}{6} = 5 \text{ s} = \frac{5}{3600} \text{ h} = \frac{1}{720} \text{ h}$.

$$\begin{aligned} M_6 &= \frac{1}{720} [v(2.5) + v(7.5) + v(12.5) + v(17.5) + v(22.5) + v(27.5)] \\ &= \frac{1}{720} (31.25 + 66 + 88 + 103.5 + 113.75 + 119.25) = \frac{1}{720} (521.75) \approx 0.725 \text{ km} \end{aligned}$$

For a very rough check on the above calculation, we can draw a line from $(0, 0)$ to $(30, 120)$ and calculate the area of the triangle: $\frac{1}{2}(30)(120) = 1800$. Divide by 3600 to get 0.5, which is clearly an underestimate, making our midpoint estimate of 0.725 seem reasonable. Of course, answers will vary due to different readings of the graph.

20. $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{2}{n} \left(5 + \frac{2i}{n}\right)^{10}$ can be interpreted as the area of the region lying under the graph of $y = (5 + x)^{10}$ on the interval

$[0, 2]$, since for $y = (5 + x)^{10}$ on $[0, 2]$ with $\Delta x = \frac{2-0}{n} = \frac{2}{n}$, $x_i = 0 + i \Delta x = \frac{2i}{n}$, and $x_i^* = x_i$, the expression for the

area is $A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(5 + \frac{2i}{n}\right)^{10} \frac{2}{n}$. Note that the answer is not unique. We could use $y = x^{10}$

on $[5, 7]$ or, in general, $y = ((5 - n) + x)^{10}$ on $[n, n + 2]$.

8. (a) Using the right endpoints to approximate $\int_3^9 f(x) dx$, we have

$$\sum_{i=1}^3 f(x_i) \Delta x = 2[f(5) + f(7) + f(9)] = 2(-0.6 + 0.9 + 1.8) = 4.2.$$

Since f is *increasing*, using *right* endpoints gives an *overestimate*.

- (b) Using the left endpoints to approximate $\int_3^9 f(x) dx$, we have

$$\sum_{i=1}^3 f(x_{i-1}) \Delta x = 2[f(3) + f(5) + f(7)] = 2(-3.4 - 0.6 + 0.9) = -6.2.$$

Since f is *increasing*, using *left* endpoints gives an *underestimate*.

- (c) Using the midpoint of each interval to approximate $\int_3^9 f(x) dx$, we have

$$\sum_{i=1}^3 f(\bar{x}_i) \Delta x = 2[f(4) + f(6) + f(8)] = 2(-2.1 + 0.3 + 1.4) = -0.8.$$

We cannot say anything about the midpoint estimate compared to the exact value of the integral.

31. (a) Think of $\int_0^2 f(x) dx$ as the area of a trapezoid with bases 1 and 3 and height 2. The area of a trapezoid is $A = \frac{1}{2}(b + B)h$,
so $\int_0^2 f(x) dx = \frac{1}{2}(1 + 3)2 = 4$.

$$\begin{aligned} \text{(b) } \int_0^5 f(x) dx &= \int_0^2 f(x) dx + \int_2^3 f(x) dx + \int_3^5 f(x) dx \\ &\quad \text{trapezoid} \quad \text{rectangle} \quad \text{triangle} \\ &= \frac{1}{2}(1 + 3)2 + 3 \cdot 1 + \frac{1}{2} \cdot 2 \cdot 3 = 4 + 3 + 3 = 10 \end{aligned}$$

(c) $\int_5^7 f(x) dx$ is the negative of the area of the triangle with base 2 and height 3. $\int_5^7 f(x) dx = -\frac{1}{2} \cdot 2 \cdot 3 = -3$.

(d) $\int_7^9 f(x) dx$ is the negative of the area of a trapezoid with bases 3 and 2 and height 2, so it equals

$$-\frac{1}{2}(B + b)h = -\frac{1}{2}(3 + 2)2 = -5. \text{ Thus,}$$

$$\int_0^9 f(x) dx = \int_0^5 f(x) dx + \int_5^7 f(x) dx + \int_7^9 f(x) dx = 10 + (-3) + (-5) = 2.$$

32. (a) $\int_0^2 g(x) dx = \frac{1}{2} \cdot 4 \cdot 2 = 4$ [area of a triangle]

(b) $\int_2^6 g(x) dx = -\frac{1}{2}\pi(2)^2 = -2\pi$ [negative of the area of a semicircle]

(c) $\int_6^7 g(x) dx = \frac{1}{2} \cdot 1 \cdot 1 = \frac{1}{2}$ [area of a triangle]

$$\int_0^7 g(x) dx = \int_0^2 g(x) dx + \int_2^6 g(x) dx + \int_6^7 g(x) dx = 4 - 2\pi + \frac{1}{2} = 4.5 - 2\pi$$

48. $F(0) = \int_2^0 f(t) dt = -\int_0^2 f(t) dt$, so $F(0)$ is negative, and similarly, so is $F(1)$. $F(3)$ and $F(4)$ are negative since they represent negatives of areas below the x -axis. Since $F(2) = \int_2^2 f(t) dt = 0$ is the only non-negative value, choice C is the largest.